

DETROIT PUBLIC LIBRARY

Vol. No. M. 2799

SUPPLEMENT: AUTHOR AND SUBJECT INDEX

FEB 11 1961

Upper Third

GENERAL INFORMATION

Current Science



Vol. 29, No. 12


DECEMBER 1960

Pages 457-500



SPECTROPHOTOMETER

Mains-operated Model PMQ II



Made by Carl Zeiss, Oberkochen, Germany

For precise measurements from 200 $m\mu$ in the ultraviolet to 1000 $m\mu$ in the near infrared region with high spectral resolution. Write for illustrated literature.


Sole Agents

ADAIR, DUTT & CO. (India) Private LTD.

CALCUTTA MADRAS SECUNDERABAD BOMBAY NEW DELHI

BAUSCH & LOMB
SPECTRONIC "505"
AUTOMATIC RECORDING SPECTROPHOTOMETER

UV-Visible Model—Double beam double grating



A revolutionary new instrument that directly records transmission, absorption, reflection and emission in the ultra-violet and visible ranges 200 m μ -700 m μ

With an exclusive electronic sensor that automatically adjusts drum speed to variations in curve complexity

Featuring B & L Certified-Precision Gratings for high resolution and minimum scattered light

Only 36" \times 22" \times 15", it's as streamlined as the universally accepted B & L Spectronic 20 Colorimeter

With a complete line of accessories including new air-cooled Hydrogen lamp

At a very competitive price for a recording Spectrophotometer




*Please write for Catalogue D-2009 to
Sole Agents in India*

MARTIN & HARRIS (PRIVATE) LTD.

(SCIENTIFIC DEPARTMENT)

SAVOY CHAMBERS, WALLACE STREET, BOMBAY 1




WILD
HEERBRUGG

* The most COMPLETE
Microscope so far
made

* Ideal for QUALITY
PRECISION
& VARIETY

WILD M 20



The M 20 Microscope with few of its attachments.

Fig. 1: With Micro Camera and Binocular Tube;

Fig. 2: Equipped as Phase Contrast Microscope;

Fig. 3: With Six-Hole Nosepiece; Filter Exchanger,
etc.;

Fig. 4: Built-in Illuminator for Kohler Illumination.

Sole Agents:
RAJ-DER-KAR & CO.

Sadhana Rayon House

Dr. D. Naoroji Road

BOMBAY-1

Telephone: 26-2304

Telegram: TECHLAB

Branch Office:

44/6 REGAL BUILDING

CONNAUGHT PLACE

NEW DELHI-1

Gansons FOR



'GS' GAS PLANTS Electrically operated
'GS' GAS PLANTS WEIGHT DRIVEN

Best suited for places without electric power supply

SUPERIOR LABORATORY EQUIPMENTS AND APPLIANCES

Ovens, Baths, Shakers, Stills, etc.

Equipments for handling and storing
RADIOACTIVE ISOTOPES

STAINLESS STEEL FABRICATIONS

Pilot Plants, Tanks, Kettles, Pressure Vessels, etc.

ORE-DRESSING PLANTS AND EQUIPMENTS

Wilfley Table, Froth Flotation Cells, etc.

GANSONS (PRIVATE) LIMITED

P.O. BOX 5576

BOMBAY 14



Model Tv 10 with metal door

AVAILABLE EX-STOCK

GERMAN 'MEMMERT' OVEN MODEL TV 10

GERMAN ROTOFIX CENTRIFUGES (Various)

WARING BLENDERS

And other Laboratory Glassware



Please contact for details:

LABORATORY FURNISHERS

DHUN MANSION, VINCENT ROAD, DADAR, BOMBAY 14

Telephone: 62761


Branch: AHMEDABAD

CRAFTSMAN

A SYMBOL OF QUALITY & SERVICE

—FOR—

LABORATORY RESEARCH
AND
INDUSTRIAL INSTRUMENTATION



Offer

- * A Complete RANGE PHOTOELECTRIC COLORIMETERS
- * OVENS, FURNACES, INCUBATORS
- * SHAKERS & WATERBATHS
- * CONTROLLERS, PYROMETERS & THERMOCOUPLES
- * DIAL THERMOMETERS



ASK FOR PRICES AND MORE DETAILS

CRAFTSMAN ELECTRONIC CORPN. (PRIVATE) LTD.

SETHNA HALL, NESBIT ROAD, MAZAGAON, BOMBAY 10

Telegram : CRAFTRONIC

Phone : 42346

PRECISION SCIENTIFIC CO., U.S.A.

SCIENTIFIC RESEARCH & PRODUCTION CONTROL
EQUIPMENT



PRECISION THERMOSTATIC ELECTRIC
THELCO DRYING OVEN

—also—

FREAS MECHANICAL & GRAVITY CONVECTION
OVENS

THELCO INCUBATORS & VACUUM OVENS

KJELDAHL DISTILLATION APPARATUS

DISTILLED WATER STILL

ETC.

FUL-KONTROL LABORATORY HEATER


with built-in auto-transformer with dial control from
0 to 750 watts. Ideal for Petroleum Laboratories

Other heating appliances as

HIGH INTENSITY TORCH

BLAST BURNERS

CROSS FIRES, ETC.



Sole Agents

GORDHANDAS DESAI PRIVATE LTD.

SIR PHEROZESHAH MEHTA ROAD, FORT, BOMBAY 1

Branches

P-7, MISSION ROW EXTENSION
CALCUTTA 1

4/2 B, ASAF ALI ROAD
NEW DELHI

22, LINGHI CHETTY STREET
MADRAS 1



BANTAM DEMINERALIZER

DOES 5 WATER PURIFICATION JOBS
EMPLOYS 5

INTERCHANGEABLE CARTRIDGES

STANDARD CARTRIDGE produces ion-free water at minimum cost. Removes 1,500 grains as NaCl (1,300 as CaCO_3).

MIXED RESIN CARTRIDGE for operations demanding better than 1,000,000 ohms resistance and neutral pH. Cartridge capacity is 1,230 grains as NaCl (1,050 as CaCO_3).

ORGANIC REMOVAL CARTRIDGE removes organics, organic liquids and gases that would pass through a demineralizer. Effective in removing chlorine. Ideal for pre-treating demineralizer feed-water, for self-purifying high purity rinse systems and other processes where organics or odors in the water are objectionable.

OXYGEN REMOVAL CARTRIDGE

was developed for cooling water loops where it is important to maintain low oxygen content to prevent corrosion. Cartridge removes one part per million of dissolved oxygen from 2,500 gallons of water, or 6.8 litres (9.7 grains, of oxygen at standard temperature and pressure.

CATION CARTRIDGE

provides (1) precious metals recovery, (2) radioactive isotope recovery at low cost, (3) also useful for removing volatile amines where heating plant steam condensate is being used as the feed-water for a Still, and (4) where close control over the pH of water is necessary, the cation cartridge in its ammonia or lithium form is effective.

In addition to these new cartridges, Barnstead has also developed several new Bantams. Among these are the BD-22, two-cartridge, two-stage laboratory unit; and the BD-10, a 1,200 G.P.H. industrial unit with disposable cartridge.

Sole Distributors

THE SCIENTIFIC INSTRUMENT COMPANY LTD.

ALLAHABAD

BOMBAY

CALCUTTA

MADRAS

NEW DELHI



only the best filter papers

You cannot be too careful about the filter papers you use.

For consistently accurate results use only GENUINE WHATMAN papers in your laboratory. They are made by modern techniques that guarantee their quality: their properties are examined and controlled at every stage of manufacture. They are preferred by leading scientists throughout the world because with Whatman you can be sure you are getting the correct results in all filtration procedures. They have no substitute. So look at the label carefully and see that it is GENUINE WHATMAN before you buy.

ensure correct results



H. REEVE ANGEL & CO. LTD

Gt. Britain: 9 Bridewell Place, London EC4


USA: 9 Bridewell Place, Clifton, New Jersey

Sole distributors of

WHATMAN FILTER PAPERS

Manufacturers W & R Balston Ltd

ZEISS MIRROR MONOCHROMATOR



Range: From 0.2 to 40 μ

An indispensable instrument for all kinds of selective measurements

VEB CARL ZEISS JENA

SOLE AGENTS

GORDHANDAS DESAI PRIVATE LTD.

SIR PHEROZESHAH MEHTA ROAD, BOMBAY 1


Branches:

P-7, MISSION ROW EXTENSION
CALCUTTA 1


4/2 B, ASAF ALI ROAD
NEW DELHI

22, LINGHI CHETTY STREET
MADRAS 1


Communications Measuring Trunk Type 4004



LCR-Precision Bridge Type 1008



Measuring Generator Type 159



P285-28/59

GERMAN ELECTRONIC MEASURING & TESTING INSTRUMENTS

Ex-Stock



- Oscilloscopes
- Signal Generators
- Precision Wavemeters
- Frequency Standards
- LCR Bridges
- Vacuum Tube Voltmeters
- Q-Meters
- Distortion Meters
- Filters, Attenuators

Precision Wavemeter Type 121



Elektrotechnik

GERMAN DEMOCRATIC REPUBLIC

Sold and serviced in India exclusively by

BLUE STAR

**BLUE STAR ENGINEERING
CO. (Bombay) PRIVATE LTD.**

KASTURI BUILDINGS
JAMSHEDJI TATA ROAD, BOMBAY I
Also at CALCUTTA, DELHI, MADRAS

PANJAB UNIVERSITY

Applications are invited for the undermentioned posts of Professors, Readers and Lecturers—Grade Rs. 800-50-1,250, Rs. 500-30-800, and Rs. 250-20-450-EB-25-650 respectively, with benefit of Provident Fund on confirmation, and allowances as admissible under the University rules. Higher starting pay in the grade depending upon qualifications and experience is admissible.


Candidates for the posts of Professors should have brilliant academic career followed by outstanding research work in the subject. They should have experience of teaching to post-graduate classes or experience of post-doctoral research of about 10 years and should be competent to guide research.

Candidates for the posts of Readers should have the Master or Doctorate or an equivalent degree in the subject concerned. They should have bright academic record, considerable research experience as indicated by published work and experience of teaching or post-doctoral research of about five years.

Candidates for the posts of Lecturers should be at least M.Sc. second class. They should have some published research work to their credit. Doctorate degree and teaching experience with Post-Graduate classes will be additional qualifications.

Reader in Chemical Engineering	1
Reader in Inorganic Chemical Technology	1
Lecturer in Chemical Engineering	1
Lecturer in Organic Chemistry	1
Professor of Chemistry (Organic Chemistry)	1
Lecturers in Chemistry.	2
(One should have specialised in Inorganic and Analytical Chemistry and the other in Bio-Chemistry)	
Lecturer in Botany	1
Lecturers in Physics	2
Reader in Pharmacology in the Department of Pharmacy	1
Lecturer in Pharmaceutical Chemistry	1

Applications, stating full particulars under the following heads: (i) Age (ii) Academic Qualifications from Matriculation or equivalent standard onwards indicating Division or class obtained in each Examination. (iii) Teaching Experience with Graduate and Post-Graduate Classes separately. (iv) Research Experience at Pre-Doctoral and Post-Doctoral Levels with appropriate details such as research papers published, research scholars guided, etc. (v) Remarks, giving additional information if any, along with copies of publications, reprints, University Certificates, testimonials, etc. should reach **The Registrar, Panjab University, Chandigarh, by January, 31, 1961.**

GLASSWARE FROM THURINGIA*A World-wide Symbol for Supreme Quality*

The history of glass-making in Thuringia can be dated back to 1597. Now, after more than 350 years, glassware for all branches of science and industry is produced here. The experience accumulated in centuries is utilised in the manufacturing of these products.

Glass for all purposes of laboratory use is manufactured in a variety of forms and shapes. Of course the pictures show only a few of the many articles offered by us. We shall be glad to send you full details of our products.

Our representatives:

PHARMA TRUST

KESHAV BAUG, 114, PRINCESS STREET
BOMBAY 2, INDIA

GLAS-KERAMIC

DEUTSCHER INNEN-UND AUSSENHANDEL
BERLIN W 8
KRONENSTRASSE 19-19A
GERMAN DEMOCRATIC REPUBLIC

Applications are invited from Indian nationals, preferably bachelors, for the post of Vegetative Propagation Adviser in North-East India for a Tea Agency House.

Applicants should possess general foreign degree in agriculture, also practical experience in plant selection and breeding essential. Work entails selection of tea bushes for the improvement of planting material and subsequent propagation by means of cuttings. Preliminary training provided. Emoluments in accordance with qualifications and experience.

Please apply clearly marked "V.P. Adviser", in own handwriting giving age and full details of degree(s), together with outline of career to date, to

DUNCAN BROTHERS & CO., LTD.

POST BOX NO. 189, G.P.O.

CALCUTTA

more
here.
ducts.
hapes,
all be



BKS
SKYREX




*High Silica low expansion heat proof resistant
glassware of best class for laboratory and industry*

Manufactured by

B.K. SHAW Industries (Private) Ltd.

P-3, B. K. PAUL AVENUE, CALCUTTA-5

Through thick and thin Pyrex



Pyrex tubing

is made from Pyrex borosilicate glass
It is therefore
heat-resistant
mechanically strong
chemically durable
—and functionally matches all other
PYREX glass apparatus

In addition, its chemical composition is now
completely free from arsenic

Sizes 2 mm to 100 mm o/diam—the largest
size-range available in borosilicate glass

Wall Thickness Standard Wall, Heavy Wall,
Extra Heavy. Also Capillary range

Length Tubing is supplied in standard lengths
of approximately 5 feet

SPECIAL problems concerning manipulated
tubing can be referred to the PYREX Service
Department, who will gladly construct special
laboratory equipment to your drawings, in
consultation (if necessary) with your scientific
and technical staff

Everyone who makes glass
apparatus needs Pyrex Tubing.
The Pyrex catalogue lists no
fewer than 80 standard lines,
all in healthy demand.
Who uses all this tubing?
Hospitals, research labs for
Government and industry,
universities, schools.
PYREX are always improving
their production methods
to attain even higher standards
of quality. This is one good
reason (among many) why
everyone who is looking
for quality glassware looks
for PYREX

PYREX

Regd. Trade Mark BRAND



Laboratory and scientific glass

Available from all leading distributors

Sole Agents: GORDHANDAS DESAI (PRIVATE) LTD.

SIR PHEROZESHAH MEHTA ROAD, FORT, BOMBAY 1

Also at: **CALCUTTA MADRAS NEW DELHI**

BOOKS from Interscience . . .

Introduction to the Theory of Quantized Fields

by N. N. BOGOLIUBOV and D. V. SHIRKOV

Authorised English Edition, Revised and Enlarged by the Authors. Translated from the Russian by G. M. Volkoff, University of British Columbia

In enlarged and revised English translation, a systematic presentation of the modern quantum field theory — from its foundations to its recent achievements. "... Will undoubtedly establish itself as one of the most important text books of the quantum theory of fields." — British Journal of Applied Physics.

1959. 736 pages. College Edition: \$ 12.50

Quantum Chemistry: Methods and Applications

by R. DAUDEL and R. LEFEBVRE, Centre de Chimie de Chimie Theorique de France, and C. MOSER, Institut du Radium

A clear presentation of the important concepts of quantum mechanics and the simple principles of the calculation of wave functions followed by a discussion of the more elaborate methods offering further progress in application.

1960. 586 pages. \$ 14.50

The Physical Chemistry of Surfaces

by A. W. ADAMSON, University of Southern California

A comprehensive physico-chemical treatment of both liquid and solid surfaces, designed as an advanced text. The theory is developed and then applied to problems of friction and lubrication; wetting, flotation and detergency; emulsions and foams; chemisorption and catalysis. Problems are included.

1960. 643 pages. \$ 12.75

Introduction to Colloid Chemistry

by KAROL J. MYSELS, University of Southern California

A long-needed text of colloid science for the undergraduate student of physical or general chemistry. The subject is treated logically and systematically; many illustrations, problems and answers supplement a clear and lucid presentation.

1959. 491 pages. College Edition: \$ 8.00

Bacteriophages

by MARK ADAMS, with chapters by E. S. Anderson, J. S. Gots, F. Jacob and E. L. Wollman.

This first comprehensive treatment in over thirty years of a subject of long-term interest is both an authoritative guide and a memorial to a pioneer in phage research.

1959. 610 pages. \$ 15.00

General Stockists:

ALLIED PUBLISHERS PRIVATE LTD.

15, Graham Road, Ballard Estate
BOMBAY 1


17, Chittaranjan Avenue
CALCUTTA 13

13/14, Asaf Ali Road
NEW DELHI 1

Available also at thirty major booksellers throughout India

Beckman Laboratory Potentiometric Recorder :

(Beckman Instruments Inc., U.S.A.)



Potentiometric Recorder

This is a low cost, push-button recorder monitoring DC signals in 10-100 mV range. It is a small and compact unit which can be operated horizontally on laboratory benches or vertically on wall. It can be used with a variety of instruments for recording of temperature, pH, frequency, heat, solution conductivity, pressure, strain speed and light intensity and can be used in the spectrophotometers UV, visible or infra-red.

Features:

Voltage span continuously variable between 10 and 100 mV full-scale multiple chart drive.

Strip chart recording and circular chart recording accessory available.

Pen travel 5 in.

Pen response 1.0 sec. full scaler.

Sensitivity 0.25% span.

Other Instruments:

pH meters, Spectrophotometer for ultra-violet, visible and infra-red regions, ultra-centrifuges, gas analysers, amino-acid analyser, paper electrophoresis apparatus, etc.

For further details contact Sole Agents:

TOSHNIWAL BROS. PRIVATE LTD.

198, JAMSHEDJI TATA ROAD, BOMBAY 1

BRANCHES:

'RIVAL', KACHERI ROAD
AJMER

11, BRITISH INDIA ST.
CALCUTTA 1

14-B/4, N.E.A., UTTARI MARG
NEW DELHI 5

ROUND TANA, MOUNT ROAD P.O.
MADRAS 2

Current Science



Vol. XXIX]

DECEMBER 1960

[No. 12

CONTENTS

	PAGE
Coherence Properties of Electromagnetic Radiation—Part II—S. PANCHARATNAM ..	457
Two-Quantum Transitions in Electron Paramagnetic Resonance ..	460
Natural Purification of Flowing Sewage—S. C. PILLAI, G. J. MOHANRAO, A. V. S. PRABHAKARA RAO, C. A. SASTRY, P. V. R. SUBRAHMANYAM AND C. V. NATARAJAN ..	461
On a Class of Asymmetrical Factorial Designs—K. KISHEN ..	465

Letters to the Editor

Ultrasonic Absorption Measurements in Some Pure Liquids—H. S. RAMA RAO AND B. RAMACHANDRA RAO ..	467	Amino-Acid Constituents of Adults of the Carpet Beetle, <i>Anthrenus vorax</i> Water- house (Dermestidae: Coleoptera)— SNEHAMOY CHATTERJI AND PRAKASH SARUP ..	479
Potential Constants of Certain Planar XY ₂ Z Molecules—P. BABU RAO AND K. SREERAMAMURTY ..	467	Utilisation of the Technique of Vegetative Propagation in Rice Breeding—R. H. RICHHARIA AND B. MISRO ..	480
Magnetism and Molecular Structure: The Estimate of Metal-Nitrogen Bond Effect in Zinc Chloride Complexes of Organic Bases—V. G. SONALKAR AND M. G. DATAR ..	468	Effect of Certain Organic Insecticides on the Yield of Crops (Part II)—D. SESHAGIRI RAO ..	480
Thermogravimetric Behaviour of Lanthanum, Thorium and Cerium Oxalates in an Atmosphere of Carbon Dioxide— O. K. SRIVASTAVA AND A. R. VASUDEVA- MURTHY ..	470	A Haematoxylin Squash Method for the Root Tips of <i>Dolichos lablab</i> Linn.—K. M. MARIMUTHU AND M. K. SUBRAMANIAM ..	482
New Colorimetric Method for the Determination of Diazinon—A. IRUDAYA- SAMY AND A. R. NATARAJAN ..	471	A Case of a Sectorial Chimera in Onion— S. L. JUNEJA AND RAM GOPAL ..	483
Propiophenone Oxime as a Reagent for the Determination of Copper—J. S. DAVE AND A. R. PATEL ..	472	A New Species of <i>Cercospora</i> on <i>Leptadenia</i> <i>reticulata</i> Wight and Arn.—N. PRASAD, R. D. SINGH AND G. C. BHATNAGAR ..	483
"Vibrated" Dropping Mercury Electrode— R. NARAYAN, V. K. VENKATESAN AND K. S. G. DOSS ..	472	Preliminary Studies on the Effect of Gibberellic Acid on Growth of Spinach (<i>Spinacia oleracea</i>)—O. S. JAUHARI, R. D. SINGH AND V. S. DIKSHIT ..	484
Heat-Resistant and Acid-Tolerant Yeast from <i>Passiflora edulis</i> Sims.—J. S. PRUTHI, B. A. S. RAO, S. A. JALEEL AND GIBDHARI LAL ..	473	The Presence of Indole-Like Compounds in Parthenocarpic Guava—G. RANGA- SWAMI AND T. T. KALIAPERUMAL ..	485
Studies on Bombay Rats: A Note on the Probable Resistance of <i>B. bengalensis</i> to Plague—P. J. DEORAS ..	475	Factors for Tillering in Sorghum Spp. (<i>S. subglabrescens</i>)—K. DIVAKARAN ..	487
Some Reactions of Phenols—J. R. MERCHANT AND V. B. DESAI ..	476	Propagation of <i>Althea rosea</i> , Benth. and Hook. (Holly Hock) by Air Layering with the Aid of Growth Regulators— D. S. LINGARAJ ..	488
Influence of Earthquake Shocks on the Arkana Gravimeter Spring—S. BALA- KRISHNA AND P. V. JOHNSON ..	476	Field Trials with Some Insecticides for the Control of Sweet Potato Weevil—K. K. NIRULA AND K. N. CHHIBBER ..	489
On the Effects of Different Concentrations of Atmospheric Oxygen on Cell-Differ- entiation in the Regenerate of the Snail, <i>Helix aspersa</i> Mull.—B. P. PANDE ..	478	Shell Disease in <i>Crassostrea gryphoides</i> (Schlothheim)—V. S. DURVE AND D. V. BAL ..	489
		Occurrence of <i>Thorea ramosissima</i> Bory in India—DALBIR SINGH ..	490

Reviews ..	491
Science Notes and News ..	498

ANNUAL REVIEW OF

MICROBIOLOGY

Vol. 14

October 1960

- Variation in Phytopathogenic Fungi, *P. R. Day*
- Microbial Nutrition, *H. C. Lichstein*
- Intramural Spread of Bacteria and Viruses in Human Populations, *R. E. O. Williams*
- Protein Synthesis in Microorganisms, *G. D. Novelli*
- Energy Metabolism in Chemolithotrophic Bacteria, *H. Lees*
- Industrial Fermentations, *S. P. Owen*
- Intracellular Symbiosis in Insects, *A. Koch*
- Nutrition, Metabolism and Pathogenicity of Microorganisms, *H. E. Adler and M. Shifrine*
- Interpretations of Immunodiffusion Tests, *A. J. Crowle*
- In Vitro* Cell-Virus Relationships Resulting in Cell Death, *D. L. Walker*
- The Genetics and Cytology of *Chlamydomonas*, *R. P. Levine and W. T. Ebersold*
- Fine Structure of Virus-Infected Cells, *H. M. Rose and C. Morgan*
- Enzyme Localization in Bacteria, *A. G. Marr*
- Arthropod-Borne Viruses of Man and Other Animals, *R. E. Kissling*
- Synchronous Division of Microorganisms, *O. H. Scherbaum*
- Gene Action, *C. Yanofsky and P. St. Lawrence*
- Immunological Speculations, *N. K. Jerne*

391 pages

NUCLEAR SCIENCE

Vol. 10

December 1960

- Neutrino Interactions, *Frederick Reines*
- Nuclear Interactions of Heavy Ions, *A. Zucker*
- Cosmic Ray Showers, *Kenneth Greisen*
- Bubble Chambers, *Hugh Bradner*
- Optics of High-Energy Beams, *Owen Chamberlain*
- Nuclear Structure Effects in Internal Conversion, *E. L. Church and J. Wenner*
- Recoil Techniques in Nuclear Reaction Studies, *B. G. Harvey*
- Labelling of Organic Compounds by Recoil Methods, *Alfred P. Wolf*
- Nucleon-Nucleon Scattering Experiments and Their Phenomenological Analysis, *M. M. MacGregor, M. J. Moravcsik and H. P. Stapp*
- Theoretical Interpretation of the Energy Levels of Light Nuclei, *I. Talmi and I. Unna*
- APPENDIX, *F. Ajzenberg-Selove and T. Lauritsen*
- Subsurface Prospecting by Nuclear Methods, *J. C. Beckerley*
- Experiments on Cosmic Rays and Related Subjects during the International Geophysical Year, *E. P. Ney*
- Cellular Radiobiology, *Tikvah Alper*
- Vertebrate Radiobiology: Metabolism of Internal Emitters, *Roy C. Thompson*
- Vertebrate Radiobiology: Late Effects, *J. B. Storer and Do. Grahn*

617 pages

each \$ 7.50 postpaid

Annual Reviews, Inc., 231 Grant Ave., Palo Alto, Calif., U.S.A.
Maruzen Company, Limited, 6 Tori-Nichome Nihonbashi, Tokyo, JAPAN

COHERENCE PROPERTIES OF ELECTROMAGNETIC RADIATION

PART II

S. PANCHARATNAM

RESONANCE RADIATION AND INDUCED EMISSION

TOWARDS the beginning of this century, R. W. Wood discovered the phenomenon of resonance radiation in sodium vapour: when the monochromatic yellow light of sodium was passed through sodium vapour, the latter in turn exhibited a yellow fluorescence of the same character, the incident radiation being rapidly absorbed on passing through the vapour. The term resonant fluorescence or resonance radiation describes the emitted light arose from the classical explanation of the phenomenon, which is worthwhile to outline in some detail—since it gives correctly the coherence properties of the secondary radiation which we shall require in later sections.

When a plane wave of light traverses a medium composed of classical dipole-oscillators the latter will be set into forced oscillation at the frequency ν of the incident wave. Because of the periodic dipole thus induced each atomic oscillator will emit radiation of the incident frequency ν —which may be observed as a feeble Rayleigh scattering transverse to the beam. However, when the frequency of the incident light coincides with the natural frequency ν_0 of the dipole-oscillators, the amplitude of the forced oscillation (and consequently the intensity of the secondary scattered radiation) should build up to large values because of the phenomenon of resonance. It must be noted that the induced dipole-oscillation will have a specific phase-relationship with the exciting light field amounting to a phase lag of $\pi/2$ at exact resonance). Thus in turn the resonance radiation from an atom should be coherent with the exciting radiation. This explains at once the apparent absence of secondary radiation in the exact forward direction, i.e., in the exact direction of the exciting beam, and the attenuation or absorption of the latter. For, it may be shown that all the secondary wavelets from a mass of atoms normal to the incident beam will coalesce to form a plane wave travelling in the same direction as the incident beam, but exactly out of phase to it. Thus, it is really the destructive interference of the resonance radiation scattered in the forward direction with the incident beam that leads to the progressive attenuation of the latter.

On the quantum theory, resonant fluorescence arises because the incident photons are of the

right energy to raise the atoms from the normal or ground state to a higher energy state, the atom then returning to the initial state with the emission of resonance radiation. The coherence of the emitted radiation with the incident, however, indicates that the two transitions, viz., to the higher energy state and back, must not be regarded as independent processes, but as part of a composite process. If we treat radiation alone classically, we may say that the oscillating dipole moment (corresponding to the double-transition) lags in phase by $\pi/2$ behind the phase of the perturbing light wave—as also happens when the atom, too, is regarded as a classical oscillator.

We have considered the atom to be initially in the lower state but the quantum picture naturally leads us to ask the following question: Could we also expect a form of resonant fluorescence when the atom is initially in the upper or excited state, and the energy of the incident photons coincides with the energy difference between the initial state and a lower energy state of the atom? The answer is yes, but the phenomenon is termed induced emission to emphasize that this radiation is induced by the presence of the light field, and thus represents something distinct from spontaneous emission: the latter is due to the natural decay of the excited atom from the upper to the lower state and would continue to exist even in the absence of an external light field. Induced emission is usually described as a transition from the initial excited state to a lower energy state under the influence of radiation of the appropriate resonant frequency, i.e., which lies within the natural line-width. However, in the author's opinion it should perhaps be considered—as in the analogous case of resonant fluorescence—as involving a double-transition, viz., to a lower energy state and back. As in the case of resonant fluorescence, the radiation from an atom due to induced emission is coherent with the light wave which stimulates it. However, an important difference in the present case is that the phase of the oscillating dipole-moment (corresponding to the double-transition of the atom) is advanced by π relative to that obtaining in normal resonant fluorescence. Thus, if we have an assembly of atoms which are—by artificial means—being kept in an excited state, then when a plane

wave of light of the appropriate resonant frequency is incident, the stimulated emission from the atoms will, in the forward direction, interfere with the original beam actually to *amplify* it—rather than to attenuate it as in the normal case of resonant fluorescence.

Such amplification has already been realised in the corresponding case of *microwaves* using the device known as the maser (see *Current Science*, 1958, Vol. 27, p. 117). But for an optical transition the difference between the energies of the upper and lower states is comparatively large, so that the problem of artificially maintaining a preponderance of atoms in an optically excited state is a difficult one. However, very recently, effects depending on light amplification by stimulated emission of radiation ('LASER' action) have been observed experimentally. We shall, in the last section, return to these experiments—which must certainly be classed among the most spectacular and fundamental experimental observations in the field of optics in recent times. Before, however, turning to stimulated emission, it would be appropriate first to deal with certain recent beautiful experiments involving resonant fluorescence which are relevant to the subject of this article.

2. INTERFERENCE EFFECTS IN CROSSED HYPERFINE LEVELS

Consider an atom with two upper energy levels separated by a small value. When irradiated with suitable radiation whose line-width is larger than this term-separation, resonance radiation of two frequencies will be emitted—corresponding respectively to transitions from the two upper levels to the ground state. The intensity of the resonance radiation will be the sum of the intensities of the radiations due to each transition, separately. If now, by the application of a magnetic field, the two upper levels are split into Zeeman components, the intensity of the emitted resonance radiation would, in general, be unaffected. However, suppose the magnetic field has a particular value such that a Zeeman sub-level of one state coincides in energy with a sub-level of the second state. Then the resonance radiations (now of identical frequency) emitted by transitions from these (coincident) sub-levels to the ground state will be coherent and in phase with one another—since both radiations should be coherent with the spectral component of the incident light responsible for exciting them. Therefore, the intensity of the resonance radiation of the frequency in question will be given by squaring the sum of the amplitudes of the radiations due

to each transition separately, and will thus be greater than the sum of the intensities due to the separate transitions. The increase in intensity of the resonance radiation will also manifest itself as an increased attenuation of the incident beam—since the latter phenomenon, as we have explained in Section 1, is due to the destructive interference of the resonance radiation with the original beam, in the forward direction. That by sweeping the magnetic field and looking for increased absorption of the exciting beam, one can determine the magnetic field strengths at which the energies of two sub-levels cross is another—this in turn giving spectroscopic information regarding the atoms. The use of such a phenomenon was described by P. A. Franken at the Conference on Coherence in a paper the title of which forms the heading to this section. Franken gave a mathematical analysis of this phenomenon from the quantum theoretical standpoint, different from the physical explanation given above by the author.

3. LIGHT BEATS IN COHERENT SCATTERING

We have mentioned above that the resonance radiations emitted by the decay of atoms from two excited states of different energies are of different mean frequencies and incoherent with one another. However, suppose an atom which is initially excited to one or other of the states is subjected to a radio-frequency field which induces transitions to the other state; then the atom may be described as being in a quantum-mechanical state which is a coherent superposition of the two states. In the subsequent decay to the ground state the radiations of the two frequencies could be expected to be coherent with one another and thus give rise to light beats. G. W. Series in the Conference at Rochester described beautiful experiments in which such a periodic modulation of the intensity of the 2,537 Å resonance radiation emitted by mercury vapour had been actually observed. The vapour is kept in a magnetic field so that the 6^3P_1 level of mercury (from which gives 2,537 Å radiation) is split into three closely spaced levels ($m_j = 1, 0, -1$), separated by equal intervals $h\nu$. A second magnetic field in a perpendicular direction oscillating at a radio-frequency ν_0 near ν , mixes the states. The photoelectric current recording the intensity of the fluorescent radiation emitted under irradiation is applied to a narrow band amplifier tuned to ν_0 , or $2\nu_0$. Light beats at the latter frequencies were thus directly detected by suitably choosing the direction of observation—the cases when the beats were absent being readily explicable as due to the

orthogonal polarisation of the two interfering radiations.

The comparative ease with which beats were observed in the above experiments is to be contrasted with the great difficulty with which beats between the Zeeman components of a spectral line were directly detected by Forrester *et al.* (see Part I). In this context, it was pointed out by the author that it becomes meaningful even when considering two quasi-monochromatic beams whose mean frequencies differ by some value ν , to talk of the coherence or otherwise of the beams—with reference to their inability to produce a beat of frequency ν . Considering the finite spectral width of each beam, each Fourier component of one beam may be paired with a corresponding Fourier component of the second beam, such that the spectral separation of the pair is ν . While the mutual interference of every such pair gives rise to a beat of the same frequency ν , the phases of the beats due to the different pairs are also identical in the case of the experiments of Series *et al.*, but not so in Forrester's experiment. This feature is vividly underlined by the remarkable fact that in the experiments of Series *et al.*, the separation in the mean frequencies of the two radiations, which may be regarded as interfering, is very much smaller than the spectral width of each radiation.

The theoretical explanation given above for the experiments of Series *et al.*, represents their original view-point as outlined very briefly in their note¹ first reporting their observation of the intensity modulation of fluorescent radiation. A feature of such an interpretation to which it seems worthwhile to draw attention is the indirect bearing it has regarding the properties of the (complex) ψ function representing the quantum-mechanical state of an atom. While normally only $|\psi|^2$, the square of the modulus, is regarded as having any physical significance, it would appear now possible to manipulate experimentally the relative phases of the ψ functions of the various states—these determining in turn the phase of the radiation emitted. It must be remembered that the above holds when the radiation field is treated classically, as is often done. However, in the Conference, Series showed that a detailed mathematical treatment in which the radiation field is quantised (following Dirac) leads directly to a periodic variation in the probability of photon emission—interpreted previously as beats.

R.F. MODULATION OF RESONANCE ABSORPTION
In 'double-resonance' experiments of the type discussed above, a modulation of the resonance

absorption of the exciting beam is to be expected due to its interference or 'beating' with the resonance radiations emitted in the exact forward direction. Such an effect had been observed in the resonance absorption due to sodium vapour by Bell and Bloom,² and the latter reported on experiments of this type. The phenomenon depended on the 'mixing' of the two ground state Zeeman sub-levels by an r.f. field of the resonant frequency. But this had to be done after first getting all the atoms into one of these Zeeman sub-levels, e.g., that for which the angular momentum vector ($J = \frac{1}{2}$) points along the direction of the magnetic field rather than against it. This was achieved by a process known as optical pumping. Modulation of optical absorption at the Larmor frequency was now observed in a circularly polarised beam transverse to the magnetic field. The experiments of Bloom *et al.*, followed a suggestion by Dehmelt,³ and in fact their interpretation of the phenomenon is of quite a different nature, i.e., not as a beat phenomenon.

5. LASER ACTION IN RUBY

In any normal monochromatic source of light the different atoms radiate incoherently. But certain recent remarkable experiments from the Bell Telephone Laboratories⁴ report what is in principle a highly monochromatic source of very large specific intensity in which all the atoms radiate coherently with one another; furthermore the radiations from the different atoms interfere constructively only along one particular direction, so that the emitted energy automatically forms a well-directed beam instead of spreading out in all directions.

The radiation in question is the well-known red fluorescent light (6,943 Å) emitted by the chromium ions in ruby. A cylindrical rod of ruby a few inches long—with the end faces optically plane-parallel and semi-silvered—was used. It was irradiated through all sides of the cylindrical surface by keeping the rod within a suitable flashlamp operated by a pulse-discharge, the fluorescent emission excited by the irradiation being examined. Under normal conditions of irradiation the fluorescent radiation from different atoms are mutually incoherent and the emission proceeds along all directions. But when the level of irradiation was gradually increased to very large values (so that the energy discharged through the lamp was of the order of 2,000 joules) the state of affairs was found to alter very suddenly. The fluorescent radiation was then found to emerge normally from the end faces as a plane parallel beam with a divergence of less than a degree.

The light as seen through the silvered ends went up by about three orders of magnitude. The spectral half-width of the 6,943 Å emission narrowed to one-thirtieth of its normal value. These experiments confirmed and extended the evidence of such behaviour first reported by Maiman.

By the absorption of light in the green portion of the spectrum the chromium ions in ruby are raised to a higher state from which they decay—practically instantaneously, for our present purposes—to an excited state 2E ; and it is the transition from this excited state back to the ground state that is accompanied by the well-known red fluorescent emission. Under normal conditions of irradiation the number of atoms in the excited state 2E will of course be very much less than in the ground state. But by increasing the intensity of irradiation, the population in the ground state can be depleted, so that actually more atoms are in the excited state. When the population inversion becomes sufficiently pronounced, the course of events become rapidly dominated by the effects due to induced emission (Section 1), and consequent coherent self-amplification of the fluorescent radiation in the ruby. One may say that each chromium ion is so engulfed by the fluorescent radiation due to the others that each ion tends to emit radiation in phase-relationship with the rest. The mutual phase-relationships which actually result in the present case is such that the resultant radiation is directed as a parallel beam along the axis. That this 'mode' of coherent emission of the assembly of ions gets

'preferred'—to the exclusion of others which one can conceive of—is due to the plane-parallel and semi-silvered ends of the cylinder. For example, a ray appreciably inclined to the axis would leave the system after a few reflections while a ray nearly parallel to the axis gets reflected back and forth, getting progressively amplified all the while (due to the population inversion in the medium).

The enormous sharpening of the spectral line width of the fluorescent emission arises mainly because the frequencies near the centre of the 'natural' line-width are more effective in stimulating emission than the rest, and are consequently amplified to a greater extent. The effect is in a sense the converse of the tendency towards self-reversal near the centre of the spectral line, observed under normal conditions where effects due to induced absorption predominate. The fact, that energy absorbed over a wide range in the green part of the spectrum is compressed to within a spectral range of 0.2 cm^{-1} , also contributes to the enormous specific intensity. In terms of a black-body emitting the same spectral density of radiation the effective temperature of the source corresponds to 10^{10} degrees kelvin!

1. Dodd, J. N., Fox, W. N., Series, G. W., and Taylor, M. J., *Proc. Phys. Soc.*, 1959, 74, 739.
2. Bell, W. E. and Bloom, A. L., *Phys. Rev.*, 1960, 107, 1559.
3. Dehmelt, H. G., *Ibid.*, 1957, 105, 1924.
4. Collins, R. J., Nelson, D. F., Schawlow, A. L., Bond, W., Garret, C. G. B. and Kaiser, W., *Phys. Rev. Letters*, 1960, 5, 303.

TWO-QUANTUM TRANSITIONS IN ELECTRON PARAMAGNETIC RESONANCE

BURGET et al., of the Institute of Nuclear

Research, Prague, report cases of two-quantum transitions in Electron Paramagnetic Resonance (EPR) spectra. The essence of two-quantum transitions consists in the simultaneous absorption of two energetically different photons or in the simultaneous absorption of one photon and the emission of another photon having a different energy (Raman process), on the assumption that the condition of the conservation of energy and the angular momentum is preserved for the system in which this transition occurs. If we apply to a system of electron spins two high-frequency fields $H_1 \cos \omega_1 t$ and $H_2 \cos \omega_2 t$, where the first is parallel to the external static field H_0 and the second is perpendicular to it, the simultaneous absorption or absorption and emission of two kinds of photons

$\hbar\omega_2$ (σ photons) and $\hbar\omega_1$ (π photons) may take place if it holds that $\hbar\omega_2 \pm \hbar\omega_1 = g\beta H_0$.

The case of emission is particularly important since the mechanism can be used to realize a maser in a system with a positive temperature while all other types of masers are based on the creation of negative temperatures in a spin system.

The apparatus used for observing these transitions consisted of an EPR spectrometer, working on a frequency $\nu_2 = 8500 \text{ MHz}$, and an autodyne spectrometer for NMR, working on frequency $\nu_1 = 13 \text{ MHz}$. The autodyne spectrometer simultaneously served as a generator of frequency ω_1 and as a detector of two-quantum transitions. The small coil of the autodyne spectrometer containing the sample, was located directly in the resonance cavity of the EPR spectrometer.—(*Czech. J. Phys.*, 1960, 10, 542)

NATURAL PURIFICATION OF FLOWING SEWAGE

S. C. PILLAI, G. J. MOHANRAO, A. V. S. PRABHAKARA RAO,
C. A. SASTRY AND P. V. R. SUBRAHMANYAM

Department of Biochemistry, Indian Institute of Science, Bangalore

AND

C. V. NATARAJAN

Formerly Deputy Director, Department of Public Health, Bangalore

[This article embodies observations made over a period of a decade and throws light on some of the fundamental principles of sanitation. The article would, therefore, be of interest to sanitary authorities throughout the world, although the set of conditions, which revealed those principles, have been provided by the peculiar method of sewage disposal particularly at Bangalore.]

It also happens that almost exactly 100 years ago, a similar but empirical observation on natural purification of flowing sewage was made in England. That was, of course, before the days of our knowledge of any microbial activity which is now known to be the most important factor governing environmental sanitation and hygiene.]

INTRODUCTION

IN the course of the discussion of a paper "On the application of town sewage to a large agricultural area, comparing its strength and dilution with the ordinary farm manurial resources; with considerations of its effects on farm profit" read before the Royal Society of Arts on March 7, 1860, one of the members of the audience, P. H. Holland, remarked: "He would in conclusion only allude to the fallacy which Mr. Sidney had put forth upon the authority of Mr. Hawksley. That gentleman, an engineer, had put forward a notion which must be astounding to every chemist. It was this, that if sewage ran away for a distance of about 10 miles, it was no longer sewage, but almost plain water. That was one of the most astounding assertions he had ever heard made".¹ Except this rather empirical statement about the behaviour of flowing sewage, which was discussed presumably against the background of the prevailing ideas of the changes in soil and sewage, there is practically no information on the natural purification of sewage in the literature. The object of this communication is to give a brief account of the conditions, particularly at Bangalore, under which some 16 million gallons of sewage now daily flows down on the outskirts of the town and purifies itself naturally without any treatment, and to indicate the operation of a principle of sanitation as evident from continued observations over a period of about ten years.

CONDITIONS AT BANGALORE

Bangalore is an inland town situated on the Deccan Plateau at a height of about 3,000 feet above sea-level, having an average annual rain-

fall of 33.3 inches, most of which occurring during May to October. The maximum temperature reached on individual days during the warmest months of April and May is 102.4° F. and the minimum temperature touched on individual days during the coolest months of December and January is 46° F. The rapidly growing town of Bangalore has now a population of about 1.4 million people and its present daily water consumption is about 17 million gallons. The water-supply is derived mostly from the Arkavati River about 22 miles west of Bangalore.

Most areas of Bangalore are provided with underground sewerage system. The bulk of the sewage is taken to three outfalls at the suburbs of the town, two of these sewage outfalls being on the southern side and the third outfall being on the south-eastern side of the town. From these outfalls the sewage is allowed to flow down in three natural channels having varying gradient, viz., 1-in-50, 1-in-100 and 1-in-800. The land surface in and around Bangalore has unusual depressions and elevations, and this topographical feature has apparently been utilised for the disposal of the town sewage since the early days of the introduction of the sewerage system in 1922.

There is no river or stream near the sewage outfalls or around the channels taking the sewage. At the various points along these channels the sewage is drawn by the farmers in the neighbourhood (over 500 families) for irrigating the land and raising crops, such as vegetables and fruits, including occasionally sugarcane and rice. There is no organised system of sewage irrigation; the farmers

improvise their own methods for taking out the sewage from the channels, when necessary.

INVESTIGATIONS AT BANGALORE

It has been observed that the turbid, foul liquid flowing in these channels becomes clear and is oxidised to an appreciable extent after its flow over a distance generally of 1 to 5 miles, depending on the gradient of the channel, and the purified effluent is used by the villagers in the vicinity for washing purposes.² In view of

these and other observations,³ a close study of the process of natural purification of flowing sewage was made, and the main results are given here. The physical features of the sewage channels at Bangalore, the zones in the channels broadly divisible in accordance with the stages of purification, the fauna and flora in them and the average results of analysis (obtained over a period of about ten years) of the sewage at different points in the channels are given in Tables I to IV. Photographs showing the condi-

TABLE I

The physical features of the sewage channels up to the regions in which the sewage are purified

	Channel having the gradient		
	1-in-50	1-in-100	1-in-800
Volume of sewage* (million gallons daily)	4	6	6
Width of the channel (feet)	4 to 12	8 to 20	4 to 10
Depth of the channel (feet)	0.3 to 1.5	0.3 to 2	0.5 to 4
Nature or type of bed	More rugged and stony at several points	Not so rugged and stony as in 1-in-50 gradient channel	Least rugged and stony
Depressions permitting sedimentation	At 0.67 mile from the outfall a considerable area of depression at certain points 1½ feet deep	Sewage stagnates at a few points	At 2.17 miles from the outfall there is a natural settling tank about ½ mile long, ½ mile wide and 3 to 4 feet deep
Dams	No dams	One dam 100 feet wide at 3.5 miles from the outfall. Three other similar dams after 6 miles from the outfall	No dams
Distance from the outfall at which sewage is completely purified (in miles)	1.29	4.75	4.46
Time taken by sewage to reach the point of complete purification (hours)	0.7 to 1	2.75 to 3.25	4 to 5
Destination of the channel	Joins 1-in-100 channel at the point 3.25 miles from the latter's origin	Joins a river about 20 miles away from the outfall	Joins a rain-fed tank 4.46 miles away from the outfall

* The sewage is mostly domestic in character and composition. About 5,000 gallons of waste water from a small tannery is introduced into the channel 1-in-50 after 0.92 mile from the outfall. About 0.3 million gallons of waste water from 3 small-sized textile mills is introduced into the channel 1-in-100 between 1.25 and 3.25 miles from the outfall. These quantities of industrial wastes did not indicate any effect on the purification of sewage.

TABLE II

The zones in the sewage channels broadly divisible in accordance with the stages of purification

Zone and stage of purification	Channel having the gradient		
	1-in-50	1-in-100	1-in-800
	Distance in miles from the outfall		
The first zone (preliminary changes leading to clarification and oxygenation)	0 to 0.67	0 to 3.50	0 to 2.71
The second zone (clarification and oxygenation)	0.67 to 0.92	3.50 to 4.00	2.71 to 3.50
The third zone (nitrification)	0.92 to 1.29	4.00 to 4.75	3.50 to 4.46
The point at which the purified effluent is used for washing purposes*	1.29	4.75 and beyond	4.46

* Pathogenic organisms have not been found at the point where the liquid is used for washing purposes. This has been ascertained with the kind assistance of the authorities of the Public Health Institute at Bangalore, to whom the authors' thanks are due.

Org.

Bacteria

Sewage
Algae

Protozoa

Rhizop.

Mastig.

Ciliop.

Rotifera

Worms

Insect Lar.

Other for.

* 1

Dominated

third zone

Distanc

in mil

from

the out

0.0

0.17

0.38

0.67

0.92

1.29

0.0

1.25

2.25

3.00

3.25

3.50

3.75

4.75

0.0

0.88

2.17

2.71

3.50

4.46

* T

† D

TABLE III
Fauna and flora in the sewage channels
(The list includes the organisms more frequently seen, and the list is by no means exhaustive)

Organisms	The first zone	The second zone	The third zone
Bacteria	.. Bacterial forms (as generally found in domestic sewage)* Occasionally <i>Salmonella typhi</i> and <i>Vibrio cholerae</i>	Aerobic forms generally	Aerobic forms generally
Sewage fungus	.. <i>Sphaerotilus</i> sp.	Not seen generally	Not seen generally
Algae	.. Not seen generally	do.	Species of <i>Oscillatoria</i> , <i>Ulothrix</i> , <i>Urospora</i> , <i>Stigeoclonium</i> and <i>Pinna-laria</i> †
Protozoa :			
Rhizopoda	.. Species, e.g., of <i>Amoeba</i> and <i>Arcella</i>	These protozoa decreased	Not seen generally
Mastigophora	.. Species, e.g., of <i>Bicosoeca</i> and <i>Euglena</i>	do.	do.
Ciliophora	.. Species, e.g., of <i>Colpoda</i> , <i>Colpidium</i> , <i>Coleps</i> , <i>Stylonychia</i> , <i>Paramecium</i> , <i>Vorticella</i> ; and occasionally species of <i>Opercularia</i> , <i>Epistylis</i> and <i>Carchesium</i>	These protozoa increased. But the species of <i>Carchesium</i> and <i>Epistylis</i> developed in strikingly large numbers forming masses of growth†	Much less growth of these protozoa
Rotifera	.. Not seen generally	Species of Rotifers	Not seen generally
Worms	.. <i>Aulophorus</i> sp.	<i>Aulophorus</i> sp.	do.
Insect Larvae	.. Mosquito (<i>Culex</i> sp.); Bloodworm (<i>Chironomus</i> sp.)	<i>Chironomus</i> sp.	Mosquito (<i>Anopheles</i> sp.).
Other forms	.. Not seen generally	Snails; fish (<i>Gambusia affinis holbrooki</i>); frogs; water hyacinth	Fish; frogs; water hyacinth

* In the first zone the bacteria predominated. † In the second zone, the protozoa, colonial Vorticellids predominated (species of *Epistylis* and *Carchesium* attached to the surfaces of stones, leaves, snails, etc.). ‡ In the third zone the algae predominated.

TABLE IV
Results of analysis of flowing sewage at different points in the channels
(Results of chemical analysis expressed as p.p.m.)

Distance in miles from the outfall	pH value	Turbidity*	3 min. permanganate value	4-hr. permanganate value	Biochemical oxygen demand	Dissolved oxygen	Ammoniacal nitrogen (N)	Albuminoid nitrogen (N)	Nitrite nitrogen (N)	Nitrate nitrogen (N)	Total bacteria (millions per ml.)
Channel having 1-in-50 gradient											
0-0	7-0	185	28-7	64-9	244	0-0†	34-2	14-0	Nil	Nil	34-0
0-17	7-4	122	17-3	36-3	167	2-1	24-9	10-4	Nil	Nil	20-0
0-38	7-5	104	13-2	24-4	128	2-7	22-8	8-5	Nil	Nil	16-0
0-67	7-5	88	9-7	17-9	99	3-6	20-5	7-0	0-02	Nil	10-0
0-92	7-8	40	5-1	11-3	32	6-0	9-7	1-94	0-20	0-37	2-0
1-29	7-8	17	2-8	9-0	15	6-7	6-3	0-56	0-44	0-61	0-003
Channel having 1-in-100 gradient											
0-0	7-3	190	25-1	50-9	219	0-0†	41-3	17-4	Nil	Nil	29-1
1-25	7-5	136	16-0	32-6	147	1-4	33-4	10-4	Nil	Nil	22-3
2-25	7-8	115	14-5	30-8	113	1-8	24-7	7-2	Nil	Nil	13-3
3-00	8-1	102	11-1	24-8	98	2-5	17-3	5-9	Nil	Nil	12-5
3-25	8-0	88	9-8	21-2	86	2-9	15-5	4-2	Trace	Trace	—
3-50	7-9	74	7-3	18-7	74	3-4	14-3	3-2	0-04	0-25	6-5
3-75	7-9	29	4-3	13-7	31	5-5	12-8	1-5	0-10	0-29	—
4-75	8-0	20	2-7	9-9	19	6-5	7-3	0-59	0-32	0-50	0-005
Channel having 1-in-800 gradient											
0-0	7-1	269	36-0	73-9	272	0-0†	54-8	18-0	Nil	Nil	35-3
0-88	7-4	188	25-2	48-9	185	2-0	37-2	10-8	Nil	Nil	27-0
2-17	7-4	106	15-5	35-0	139	2-7	28-9	8-0	Nil	Nil	15-0
2-71	7-6	72	8-9	21-5	99	3-7	18-4	5-4	0-04	Nil	10-0
3-50	7-8	47	5-3	14-3	58	6-8	12-8	0-93	0-50	0-10	0-05
4-46	8-0	23	3-0	11-1	18	6-8	9-1	0-45	0-43	0-57	0-01

* Turbidity : Values obtained by using Klett-Summerson photoelectric colorimeter with 420 filter.

† Dissolved oxygen : Occasionally the sewage samples at the outfalls contained negligible amounts of oxygen.

tions at four points in the channel having 1-in-50 gradient are given in Figs. 1 to 4.

the sewage (at 10-25% level, by volume) and the mixture gently shaken or into which air was



FIG. 1



FIG. 2



FIG. 3



FIG. 4

FIGS. 1-4. Fig. 1. One of the sewage outfalls at Bangalore: the sewage flows down in an open channel having 1-in-50 gradient. Fig. 2. At 0.67 mile from the outfall, the sewage gushes down a rugged stony area. Fig. 3. Photo micrograph of *Carchesium* sp. (x about 89) found in fluffy masses down the stony area. Fig. 4. The clear, purified effluent (at 1.29 miles from the outfall) being used by washermen.

In the light of the evidence accumulating at Bangalore the more important factors influencing purification of the flowing sewage include: (1) adequate agitation or turbulence of the sewage and other conditions in the channel, which facilitate the dissolution of oxygen to the extent of about 3.5 p.p.m., and (2) the consequent development in large numbers of ciliate protozoa notably of the species of *Carchesium* and *Epistylis* which are always found in activated sludge.⁴ When these protozoan colonies were taken out, washed and introduced into

bubbled for 1½-6 hours (depending on the number of the organisms), it was observed that the sewage was clarified and oxygenated almost to the same extent as under the natural conditions in the channels. The flocculating activity of the protozoa and the clarification of the sewage also seemed to explain the relatively high nitrogen contents of the soils under the flowing sewage in the zone of clarification and to bring about nitrification and other changes e.g., rapid removal of amino acids^{5,6} from the sewage, in the succeeding stages of purification.

the channels. The quality of the final effluents from these channels was similar to that from the activated sludge process.

OBSERVATIONS MADE AT OTHER PLACES IN INDIA

The evidence collected on the flowing sewage at the Mysore sewage farm, Shimoga, Bhadravati and Madurai⁷ also indicated that the extent of agitation of the sewage during its flow is a basic factor in the process of natural purification as it influences the initial oxygenation of the sewage, development of the protozoa and the consequent changes leading to purification. When, however, sewage flowed down in a more or less contour channel, it was not oxygenated or purified to any appreciable extent even after its flow over a distance of 5 miles.⁷

SIGNIFICANCE OF THE OBSERVATIONS

Natural purification of flowing sewage is thus essentially an aerobic process and, under the most favourable conditions, it would proceed rapidly, as observed in the channel having 1-in-10 gradient, and give results attainable only

by the activated sludge process. The above observations are of scientific interest as well as of practical importance as they not only relate to a sanitary principle in Nature and its bearing particularly on the modern methods of sewage disposal but indicate the possibility of increasing the efficiency of the activated sludge process and other methods of aerobic treatment of sewage.

1. Mechi, A., *J. Soc. Arts*, 1860, **8**, 261.
2. Pillai, S. C., Monanrao, G. J., Prabhakara Rao, A. V. S., Natarajan, C. V. and Madappa, I. B., *Indian Med. Gaz.*, 1953, **88**, 507.
3. Pillai, S. C., *Ibid.*, 1952, **87**, 411.
4. ———, *Curr. Sci.*, 1941, **10**, 84; 1942, **11**, 437; 1949, **18**, 382.
5. Ananteswara Sastry, C., Subrahmanyam, P. V. R. and Pillai, S. C., *Sewage and Industrial Wastes*, 1958, **30**, 1241.
6. Subrahmanyam, P. V. R., Anandeswara Sastry, C., Prabhakara Rao, A. V. S. and Pillai, S. C., *J. Water Pollution Control Federation*, 1960, **32**, 344.
7. Pillai, S. C., *Investigations on Sewage Farming*, Indian Council of Agricultural Research, 1955.

ON A CLASS OF ASYMMETRICAL FACTORIAL DESIGNS

K. KISHEN

Chief Statistician to Government, U.P., Department of Agriculture, Lucknow

KISHEN AND SRIVASTAVA^{1,2} have developed general methods for the construction of asymmetrical factorial designs. One of the methods given by them is for the construction of the class of $q \times 2^2$ factorial designs—the treatments being all combinations of the three factors A (0, 1, 2, ..., $q-1$), B (0, 1) and C (0, 1)—in $2b$ blocks of $2q$ plots each by use of the associated BIB design $v = q, b, k, r, \lambda$. They have illustrated the method by constructing the design 7×2^2 in 14 blocks of 14 plots each by use of the associated BIB design $v = 7, b = 7, k = 3, r = 3, \lambda = 1$ and have shown that in this design the loss of information on BC

property of balanced designs. In general, for the $q \times 2^2$ design in $2b$ blocks of $2q$ plots each, constructed from the associated BIB design $v = q, b, k = t, r, \lambda$, it can be shown that

$$L(BC) = \frac{(q-2t)^2}{q^2} \quad (3)$$

and

$$L(ABC) = \frac{4t(q-t)}{q^2(q-1)}, \quad (4)$$

the total loss of information being, as before, unity.

It would be seen from (3) that when q is odd, the minimum loss of information on BC would be $1/q^2$ for $t = (q-1)/2$. Also, when q is even, this minimum loss would be 0 for $t = q/2$. These would be called optimum designs and can be constructed when the associated BIB designs exist. In the latter case, i.e., when q is even, optimum designs can be constructed with only b blocks of $2q$ plots each, thus reducing the number of blocks otherwise required by half. In this note, only designs for $q \leq 19$ have been discussed as optimum or near optimum designs can be constructed only in these

channel
any area.
ny area.
on the
ved the
ed almost
al condi
g activ
n of the
relative
under
ation an
change
from the
purifica

$$L(BC) = \frac{1}{49} \quad (1)$$

and that on each of the 6 degrees of freedom of ABC by

$$L(ABC) = \frac{8}{49}, \quad (2)$$

that the total loss of information on the partially confounded degrees of freedom for BC and ABC in this design is unity, which is a

cases owing to the existence of the associated BIB designs with the desired parameters, considering only BIB designs with $r \leq 10$.

2. *Designs for q Odd.*—We may first discuss designs for q odd. It is known that for $q = 5$ and 9, BIB designs with $k = 2$ and 4, $b = 10$ and 18 respectively exist. In the case of 5×2^2 and 9×2^2 designs, it is possible to construct optimum balanced designs with only 10 and 18 blocks respectively instead of 20 and 36 blocks respectively required by use of the associated BIB designs. For the case 5×2^2 , Shah³ has constructed such a design. A similar design for 9×2^2 has been constructed by first obtaining the design in 18 replications from the associated BIB design $v = 9$, $b = 18$, $k = 4$, $r = 8$, $\lambda = 3$ by writing down the set " X_0, X_1 " in the BIB pattern (where X_0 denotes the combinations 00 and 11, and X_1 the combinations 01 and 10 of factors B and C), filling in the remaining places by the set " X_1, X_0 ", and then taking from 9 of the resulting replicates only those blocks which have 4 X_0 's and 5 X_1 's and from the remaining 9 replicates only those blocks which have 5 X_0 's and 4 X_1 's. In this design

$$L(BC) = \frac{1}{81} \quad (5)$$

and

$$L(ABC) = \frac{10}{81}, \quad (6)$$

the total loss of information being unity, as required.

For $q = 7, 11, 15$ and 19, BIB designs with $b = 7, 11, 15$ and 19 and $k = 3, 5, 7, 9$ respectively exist. Consequently, for the cases 7×2^2 , 11×2^2 , 15×2^2 and 19×2^2 , optimum designs in 14, 22, 30 and 38 blocks of 14, 22, 30 and 38 plots each respectively have been obtained in which the loss of information on BC is $1/q^2$ and on each of the $q - 1$ degrees of freedom of ABC is $(q + 1)/q^2$, where $q = 7, 11, 15$ and 19 respectively.

3. *Designs for q Even.*—We may now discuss designs when q is even. In this case, as remarked above, balanced designs in b blocks of $2q$ plots each can be readily constructed by use of the associated BIB designs with parameters $v = q$, b , $k = q/2$, r , λ for values of q for which these exist. Here the X_0 's alone are written in the BIB pattern instead of the set " X_0, X_1 " for q odd, and the remaining $q/2$ places in each of the b blocks are filled in by X_1 's. Obviously, in these designs, BC is un-

confounded. As BIB designs with the above parameters exist only for $q = 4, 6, 8$ and 10 (even values of $q < 19$ ($r \leq 10$)), the required type of optimum design exist only in the cases 4×2^2 , 6×2^2 , 8×2^2 and 10×2^2 . In these designs,

$$L(BC) = 0$$

and

$$L(ABC) = \frac{1}{q-1} \quad (q=4, 6, 8, 10) \quad (8)$$

so that the total loss of information is, as before, unity, which is a property of balanced designs.

4. *Near optimum designs.*—For $q = 13$, no BIB design with $k = 6$, exists. No optimum design with $L(BC) = 1/36$ can, therefore, be constructed in the case of the 13×2^2 design. However, a BIB design with $k = 4$, $b = 13$ does exist, which enables the construction of a 13×2^2 design in 26 blocks of 26 plots each in which

$$L(BC) = \frac{25}{169} \quad (9)$$

and

$$L(ABC) = \frac{12}{169}. \quad (10)$$

This is the most efficient design available in this case.

For $q = 16$, no BIB design with $k = 8$ exists. Consequently, no optimum design of the type discussed in para 3 can be constructed in the case of the 16×2^2 design. However, a BIB design with parameters $v = 16$, $b = 16$, $k = 6$, $r = 6$, $\lambda = 2$ exists, which gives a design in 32 blocks of 32 plots each by the method indicated in para 2 for q odd. In this design,

$$L(BC) = \frac{1}{16} \quad (11)$$

and

$$L(ABC) = \frac{1}{16}. \quad (12)$$

This is the most efficient design available in this case.

For full details regarding the construction and analysis of the designs discussed in this note, the interested reader is referred to the author's paper on the subject to be published elsewhere.

1. Kishen, K. and Srivastava, J. N., *Curr. Sci.*, 1969, **28**, 98-100.
2. — and —, *J. Ind. Soc. Agric. Stat.*, 1960, **11** (1 & 2), 73-110.
3. Shah, B. V., *Biometrics*, 1960, **16**, 115-18.

LETTERS TO THE EDITOR

ULTRASONIC ABSORPTION MEASUREMENTS IN SOME PURE LIQUIDS

In recent years considerable attention is being given to the study of ultrasonic absorption in liquids because of the anomaly that the observed absorption coefficients are usually greater than the theoretical values. The authors have recently developed a technique for the study of absorption in corrosive liquids by using the diffraction method and the results of absorption measurements carried out for some esters and a corrosive liquid using this technique are presented in this communication.

The ultrasonic absorption is measured by the well-known optical diffraction method.¹⁻³ In this method a parallel beam of monochromatic radiation of constant intensity is passed through the experimental medium which is itself traversed by plane progressive ultrasonic waves in a direction perpendicular to the direction of light propagation. This results in the diffraction of the emergent light, and a measurement of the diffracted light intensity of the first order which is proportional to the sound intensity, for low values of sound intensity, enables the estimation of the ultrasonic absorption coefficient. In the present investigation a sodium vapour lamp driven by a constant voltage transformer is used as the source. The crystal holder is a pyrex glass sheet with a deep circular depression in the centre, which serves simultaneously as the lid of the ultrasonic cell. The bottom surface of this depression will be in contact with the liquid in the cell. The top surface of the lid is silvered and the piezo crystal rests in the depression with a film of oil to give acoustic contact. An electrode pressing on the top silvered surface of the crystal and another on the silvered surface of the lid will serve as the electrodes for exciting the crystal. The diffracted light intensity of the first order is measured accurately by means of a sensitive Photovolt multiplier photometer. An X-cut quartz crystal driven at one of its odd harmonics by a low power, variable frequency Hartley oscillator generated the ultrasonic waves. The frequency is maintained constant using a heterodyne wavemeter of high accuracy. The specially designed all-glass crystal holder was necessary to isolate the corrosive liquid from the silvered quartz crystal. Measurements were carried out at

high frequencies of 25 and 35 Mc./sec. such that progressive waves can be easily set up in a small quantity of liquid in the cell. Temperatures of the liquids are maintained constant to within 0.1° C. The values of α/f^2 thus obtained are presented in Table I.

TABLE I

Liquid	Temp. in °C.	$(\alpha/f^2) \times 10^{17}$ in nepers/ sec. ² cm. ⁻¹		
		Experimental values		Theoretical values
		25 Mc./ sec.	35 Mc./ sec.	
Phosphorus oxy- chloride	31.7	199	212	..
Methyl Benzoate	32.7	62.7	61.3	16
Ethyl Benzoate ..	35.0	68.0	49	21
Benzyl Benzoate	40.5	158	132	45
Benzyl formate ..	32.9	..	53.3	14
Benzyl acetate ..	34.2	..	63	18

It is evident from Table I that the observed ultrasonic absorption is considerably higher than the theoretical value, a common feature which has been reported in many liquids. The apparent discrepancy in the observed values of α/f^2 for the two different frequencies in some of the liquids studied is characteristic of the particular liquids and warrants a detailed study of the absorption variation over a wide range of frequencies.

The authors wish to express their thanks to the Council of Scientific and Industrial Research for the financial aid in connection with this project.

Ultrasonic Labs., H. S. RAMA RAO.
Physics Department, B. RAMACHANDRA RAO.
Andhra University,
Waltair, November 3, 1960.

1. Willard, G. W., *J.A.S.A.*, 1947, 12, 48.
2. Burton, C. J., *Ibid.*, 1948, 20, 186.
3. Krishna Murty, M., *Proc. Ind. Acad. Sci.*, 1956, 43, 106.

POTENTIAL CONSTANTS OF CERTAIN PLANAR XY₂Z MOLECULES

Using Wilson's¹ FG matrix method and employing the most general force field, the force constants of CH₂O, CD₂O, CF₂O, CCl₂O, BF₂Cl and BBr₂Cl are evaluated. The usual notation is employed in listing the constants in Table I.

TABLE I
 f is given in units of 10^5 dynes/cm. $p = 2(f_a - 2f_{aa}) + (f_a + f_{aa})$

Molecule	f_D	f_a	f_{aa}	f_{Da}	p	$f_a - f_{aa}$	$f_{Da} - f'_{Da}$	f_τ
CH ₂ O ..	12.746	4.373	0.033	0.280	1.189	0.704	0.050	0.996
CD ₂ O ..	13.347	4.882	1.934	1.258	3.313	1.029	0.200	0.655
CF ₂ O ..	11.836	2.471	1.049	-0.080	1.143	0.204	-0.101	0.117
CCl ₂ O ..	3.375	4.405	1.537	0.803	2.331	1.237	-0.700	0.304
B ¹¹ F ₂ Cl ..	4.801	2.276	0.572	1.291	0.549	0.309	-0.016	0.108

TABLE II

Molecule	Authors	f_a	f_D	f_{Da}	f
CH ₂ O ..	Venkateswarlu and Sundaram	4.338	11.420	-0.434	0.901
	Present work	4.373	12.746	0.280	0.996
CF ₂ O ..	Lovell <i>et al.</i>	4.838	14.387	..	0.334
	Venkateswarlu and Sundaram	8.381	12.320	1.380	0.441
	Present work	4.882	13.347	1.258	0.655
CCl ₂ O ..	Venkateswarlu and Sundaram	2.369	12.860	0.943	0.119
	Present work	2.471	11.836	-0.08	0.117

TABLE III

Molecule	d (Å) (Calculated) (Badger's rule)	d (Å) (Experimental)	D (Å) (Calculated) (Badger's rule)	D (Å) (Experimental)
CH ₂ O ..	1.09	1.07	1.21	1.23
CD ₂ O ..	1.36	1.32	1.20	1.17
CF ₂ O ..	1.79	1.74	1.22	1.18
CCl ₂ O ..	1.43	1.29	1.69	1.73
BF ₂ Cl ..	1.66	1.87	1.67	1.73

D , d represent the X-Z and X-Y bond distances; A , a the Y-X-Y and Y-X-Z interbond angles and τ the out-of-plane displacement of X-Z bond with respect to the XY₂ plane.

Table II shows a comparison of the force constants with those of earlier workers,^{2,3} who used a less general force field.

Interatomic distances are calculated using Badger's empirical rule⁴ and the force constants. These are listed in Table III along with the experimental values.

The vibrational frequencies of the molecules recalculated using the above force constants are in good agreement with the observed frequencies.

Details are being published elsewhere.

Dept. of Physics, P. BABU RAO.
 S.V. University College, K. SREERAMAMURTY.
 Tirupati, November 1, 1960.

1. Wilson, E. B. Jr., *J. Chem. Phys.*, 1939, 7, 1041; 1941, 9, 76.
2. Lovell, R. J., Stephenson, C. V. and Jones, E. A., *Ibid.*, 1954, 22, 1953.
3. Venkateswarlu, K. and Sundaram, S., *Ibid.*, 1957, 202.
4. Badger, R. M., *Ibid.*, 1934, 2, 128; 1935, 3, 710.

MAGNETISM AND MOLECULAR STRUCTURE: THE ESTIMATE OF METAL-NITROGEN BOND EFFECT IN ZINC CHLORIDE COMPLEXES OF ORGANIC BASES

VARIOUS attempts have been made to correlate the basicity of the donor atom in a complexing molecule with its ability to form a metal complex. Among the various methods which have been employed to get the idea of the metal ligand bond strength, magnetic susceptibility also has been of use. In the present investigation an attempt has been made to estimate the relative strengths of metal-nitrogen bond for different addenda in the complexes of zinc with pyridine, quinoline, and aniline and its derivatives by comparing their molar susceptibilities with that of zinc diammine.

The pyridine and quinoline complexes of zinc were prepared by following the methods suggested by Lang¹ and Borsbach² respectively. Complexes of aniline and its derivatives were prepared by the methods suggested by Lachowicz and Bandrowski³ and Tombeck.⁴ The purity of these complexes was ascertained by estimating the halogen contents in them.

Magnetic susceptibilities of the complexes and their components were measured by a modified form of Gouy's balance described by Prasad *et al.*⁵ These results are given in Table I A in which χ and χ_m denote respectively the specific and molar susceptibilities expressed in -1×10^6 g.s. units.

the ammonia molecule. This difference (denoted by $\Delta \chi_m$) should be equal to the difference between the molar susceptibilities of the pure ligands (denoted by $\Delta \chi_m'$) assuming the strength of the bond between metal and nitrogen in either complex to be the same. Any difference in the two sets of values may be regarded

TABLE I A

Name of the complex	χ	χ_m	Name of ligand	χ_m
Dipyridine zinc chloride	.. 0.489	143.9	Pyridine	46.92
Diquinoline zinc chloride	.. 0.547	215.9	Quinoline	87.52
(i) Dianiline zinc chloride-dihydrate (<i>ortho</i>)	.. 0.545	(i) 195.5	Aniline	62.55
(ii) Anhydrous	(ii) 174.56
Ditoluidine zinc chloride-dihydrate (<i>ortho</i>)	.. 0.577	(i) 222.8	<i>o</i> -Toluidine	74.54
		(ii) 201.86 (anhydrous)
Ditoluidine zinc chloride (<i>meta</i>)	.. 0.583	193.9	<i>m</i> -Toluidine	74.40
Ditoluidine zinc chloride (<i>para</i>)	.. 0.554	(i) 213.9	<i>p</i> -Toluidine	72.58
		(ii) 192.96 (anhydrous)
Di-xylylene zinc chloride-dihydrate (<i>meta</i>)	.. 0.560	(i) 232.3	<i>m</i> -Xylylene	86.90
		(ii) 211.36 (anhydrous)
Diammine zinc chloride	98.77	Ammonia (liquor)	19.00

TABLE I B

Complex	χ_m	$\Delta \chi_m$	Ligand	χ_m'	$\Delta \chi_m'$	$\frac{\Delta \chi_m - \Delta \chi_m'}{2} = \lambda$ per Zn-N bond
$\text{ZnCl}_2 \cdot 2\text{C}_5\text{H}_5\text{N}$.. 143.9	45.13	$2\text{C}_5\text{H}_5\text{N}$	93.84	55.84	- 5.35
$\text{ZnCl}_2 \cdot 2\text{NH}_3$.. 98.77		2NH_3	38.00		
$\text{ZnCl}_2 \cdot 2\text{C}_6\text{H}_7\text{N}$.. 215.9	117.13	$2\text{C}_6\text{H}_7\text{N}$	175.04	137.04	- 9.95
$\text{ZnCl}_2 \cdot 2\text{NH}_3$.. 98.77		2NH_3	38.00		
$\text{ZnCl}_2 \cdot 2\text{C}_6\text{H}_5\text{NH}_2$.. 174.56	75.79	$2\text{C}_6\text{H}_5\text{NH}_2$	125.10	87.10	- 5.65
$\text{ZnCl}_2 \cdot 2\text{NH}_3$.. 98.77		2NH_3	38.00		
$\text{ZnCl}_2 \cdot 2\text{C}_7\text{H}_7\text{NH}_2$ (<i>ortho</i>)	201.86	103.00	$2\text{C}_7\text{H}_7\text{NH}_2$ (<i>ortho</i>)	149.08	111.08	- 3.99
$\text{ZnCl}_2 \cdot 2\text{NH}_3$.. 98.77		2NH_3	38.00		
$\text{ZnCl}_2 \cdot 2\text{C}_7\text{H}_7\text{NH}_2$ (<i>meta</i>)	.. 193.9	25.13	$2\text{C}_7\text{H}_7\text{NH}_2$ (<i>meta</i>)	148.80	110.80	- 7.83
$\text{ZnCl}_2 \cdot 2\text{NH}_3$.. 98.77		2NH_3	38.00		
$\text{ZnCl}_2 \cdot 2\text{C}_7\text{H}_7\text{NH}_2$ (<i>para</i>)	.. 192.96	94.17	$2\text{C}_7\text{H}_7\text{NH}_2$ (<i>para</i>)	145.16	107.16	- 6.48
$\text{ZnCl}_2 \cdot 2\text{NH}_3$.. 98.77		2NH_3	38.00		
$\text{ZnCl}_2 \cdot 2\text{C}_8\text{H}_9\text{NH}_2$.. 211.36	112.59	$2\text{C}_8\text{H}_9\text{NH}_2$	173.80	135.80	- 11.80
$\text{ZnCl}_2 \cdot 2\text{NH}_3$.. 98.77		2NH_3	38.00		

The complexes studied in this investigation are analogous to the diammine of zinc, susceptibility of which has been measured by Khopkar⁶ in this laboratory. If it is assumed that ammonia molecule in zinc diammine is substituted by the ligand molecule, the difference in the molar susceptibility of the two complexes should represent the difference in the susceptibility of the ligands, viz., the organic base and

as depression due to the bond strength between the metal and nitrogen. The calculated values of $\Delta \chi_m$ and $\Delta \chi_m'$ are given in columns 3 and 6 respectively of Table I B. The differences in $\Delta \chi_m$ and $\Delta \chi_m'$ (denoted as λ) are given in the last column of the same table.

The molar susceptibilities of anhydrous complexes were obtained by subtracting the calculated value of the susceptibility of water mole-

and cerous oxalate nona hydrate $\text{Ce}_2(\text{C}_2\text{O}_4)_3 \times 9\text{H}_2\text{O}$. Thorium oxalate begins to lose water at 60° in air (curve B_1) and this continues till about 140° C. when a tri-hydrate is obtained. The weight remains constant till 200° C. where there is again a slight loss in weight. In the range 240° – 300° the sample was found to correspond to a dihydrate. (The dihydrate could be prepared in a separate experiment by heating the hepta hydrate at 240° C. for 2 hours.) The dihydrate loses water at 300° C. and passes through a mono-hydrate stage at 340° beyond which the decomposition becomes rapid. The weight of the residue becomes constant at 600° C. and remains the same till 640° C. and the composition of the final product corresponds to ThO_2 .

In an atmosphere of carbon dioxide the dehydration process of thorium oxalate is similar to that in air (curve B_2) till about 300° C. and further weight loss is very much slowed down. The thorium level is reached only beyond 600° C. The residue was dark in colour containing some elemental carbon along with thorium.

The thermogravimetric behaviour of cerous oxalate is somewhat different from the above two salts. The decomposition of cerous oxalate is very much slowed down in carbon dioxide (curves C_1, C_2). If the oxalate is heated in air all the water is lost at about 220° C. and decomposition is quite rapid at the same time. The curve C_1 is very steep between 240° and 260° C. and becomes horizontal at 320° C. The residue left behind is ceric-oxide (CeO_2). While in carbon dioxide the decomposition becomes rapid only after 320° and the curve reaches horizontal level only beyond 440° and the weight of the sample remains constant till about 440° . The residue left was ceric oxide (CeO_2) and no elemental carbon was found to be present along with it.

The thermogravimetric results in air can be compared with the results of earlier workers.³⁻⁹ The results presented here indicate that anhydrous oxalates are very unstable and it is very difficult to make out the two stages corresponding to complete dehydration and decomposition, when they are heated in air. In an atmosphere of carbon dioxide the decomposition can be delayed even after dehydration stage is reached. The delay in decomposition may perhaps be attributed to the delay in nucleation, dissociation of the carbonate and the oxidation of the oxalate. Further work is in progress to elucidate this phenomenon.

The authors wish to express their grateful thanks to Prof. M. R. A. Rao for his keen interest in the work.

Dept. of Inorganic and O. K. SRIVASTAVA.
Physical Chemistry, A. R. VASUDEVAMURTHY.
Indian Institute of Science,
Bangalore-12, October 12, 1960.

1. Fernelius, W. C., *Inorganic Synthesis*, McGraw-Hill Book Company, Inc., 1946, 2, 59.
2. Vasudevamurthy, A. R., Bharadwaj, D. K. and Mallaya, R. M., *Chem. and Ind.*, 1956, 300.
3. Backer, H. J. and Klaassens, K. H., *Z. anal. Chem.*, 1930, 81, 104.
4. d'Eye, R. W. N. and Sellman, P. G., *J. Inorg. Nucl. Chem. G.B.*, 1955, 1, 143.
5. Takayuki Somuya and Shizo Hirano, *J. Soc. Chem. Ind. Japan*, 1939, 34, suppl. binding No. 11, 459.
6. Wendlandt, W. W., *Anal. Chem.*, 1958, 30, 58.
7. Chase, W. S., *J. Am. Chem. Soc.*, 1917, 39, 576.
8. Caro, P. and Lorriers, J., *J. Rech. C.N.R.S.*, 1957, 39, 107.
9. Beckett and Winfield, *Austr. J. Sci. Res.*, 1951, 4, 644.
10. Srivastava, O. K. and Vasudevamurthy, A. R., *J. Sci. Ind. Res. (India)*, 1960, 19B, 364.

A NEW COLORIMETRIC METHOD FOR THE DETERMINATION OF DIAZINON

SEVERAL enzymic,¹⁻⁴ bioassay⁵ and ultra-violet spectrophotometric methods^{6,7} are available for the determination of diazinon. Colorimetrically diazinon is determined either by the method of Suter *et al.*,⁸ via the methylene-blue reaction (λ_{max} 670 m μ), or by the determination of total phosphorus by the method of Holman⁹ via the molybdenum-blue reaction (λ_{max} 650 m μ). These authors⁸ have also used turbidimetric tests with heteropoly acids and infra-red spectroscopy for the detection of small amounts of diazinon. The present note reports a new colorimetric method for the determination of diazinon in biological materials, in residue analysis, etc., by the application of the well-known Janovsky reaction.

An aliquot extract of the sample to be analysed containing about 5–50 μg . of the insecticide is cleaned up by chromatographic, partition separation or other suitable method. The insecticide is then nitrated and the resulting nitro-compound, after extraction, is reacted with methyl ethyl ketone in presence of strong alkali and the transmittancy of the red colour formed is measured at 502 m μ . The minimum detection level is 5 μg . of diazinon.

Sulphur and fungicides of the thiocarbamate type which interfere in the method of Suter *et al.*,⁸ and arsenic, silica and other phosphates, which interfere in the Holman method,⁹ do not interfere in the new colorimetric method. Interference is caused by compounds which readily get nitrated and especially when the product is a di- or polynitro derivative with two

nitro groups formed in the metaposition to each other. But this interference can be eliminated by suitable separation techniques prior to the nitration step.

The detailed paper will appear elsewhere.

State Forensic Sciences Lab., A. IRUDAYASAMY.
Madras-3, A. R. NATARAJAN.
September 21, 1960.

1. Giang, P. A. and Hall, S. A., *Anal. Chem.*, 1951, **23**, 1830.
2. Frawley, J. P., Laug, E. P. and Fitzhugh, O. G., *J. Assoc. Offic. Agr. Chemists*, 1952, **35**, 745.
3. Cook, J. W., *Ibid.*, 1954, **37**, 561.
4. Fallscheer, H. O. and Cook, J. W., *Ibid.*, 1956, **39**, 691.
5. Terriere, L. C. and Kiigemagi, U., *Simplified Bioassay for Insecticide Residues*. Presented at the 129th National Am. Chem. Soc. Meeting, April 1956.
6. Harris, H. J., Geigy Co., *A Tentative Ultraviolet Method for Analysis of Diazinon in Spray Residues*, September 1953.
7. Blinn, R. C. and Gunther, F. A., *J. Agr. Food Chem.*, 1955, **3**, 1013.
8. Suter, R., Delly, R. and Meyer, R., *Z. anal. Chem.*, 1955, **147**, 173.
9. Holman, W. I. M., *Biochem. J.*, 1943, **87**, 256.

RESPROIOPHENONE OXIME AS A REAGENT FOR THE DETERMINATION OF COPPER

RESACETOPHENONE oxime has been used for the estimation of copper.^{1,2} In this note the use of resproiophenone oxime as a reagent for the estimation of copper is described.

Resproiophenone³ was converted into its oxime by treatment with hydroxylamine hydrochloride. The oxime was recrystallised from ethyl alcohol, m.p. 189° C. (Lit. 186-87° C.).

10.0 ml. of the copper chloride solution was neutralised with 2 N sodium hydroxide solution and then acidified with 2 N acetic acid solution. The solution was diluted to about 200 ml. and heated to boiling. An excess of 1% solution of the reagent was then added dropwise with constant stirring and the solution along with the precipitate was heated to boiling. The brown precipitates obtained were filtered through sintered glass crucible and washed with hot water containing acetic acid till the filtrate gave no colour with ferric chloride solution. The precipitates were dried at 110°-120° C.

The copper content of the complex was determined by the decomposition method on the basis of the formula $\text{Cu}(\text{C}_9\text{H}_{10}\text{O}_2\text{N})_2$ (Cu Found: 14.97, 14.98, 15.08; Calc. 15.00 m.p. 284° C.). Further work is in progress.

The authors are thankful to Professor S. M. Sethna for his interest in the work and Dr. A. M. Talati for useful discussion.

Chemistry Department,
Faculty of Science,
M.S. University of Baroda,
Baroda, September 2, 1960.

J. S. DAVE.
A. R. PATEL.

1. Raja and Neelakantam, K., *Curr. Sci. (India)* 1950, **19**, 383.
2. Bhatki, K. S. and Kabadi, M. B., *J. Univ. Baroda* 1955, **24** (3), 51.
3. Brewster, C. M. and Harris, J. C., *J. Am. Chem. Soc.*, 1930, **52**, 4669.

"VIBRATED" DROPPING MERCURY ELECTRODE

THE behaviour of dropping mercury electrode (DME) in an external alternating field has been recently studied by Hideo Imai *et al.*¹ They found that the polarographic limiting current in the presence of the alternating field increased beyond the diffusion controlled one. At lower values of the alternating field, the limiting current coincides with the diffusion current at the potential of the electro-capillary maximum (ECM). However, at sufficient high values of the field, the limiting current does not coincide with the diffusion current even at the potential of ECM. This technique, which they have called 'vibrated' dropping mercury electrode (VDME), has been used to determine the potential of the ECM. The above workers have ascribed the increased limiting current to the streaming of solution on and near the DME caused by the vibration of the DME in alternating field (i.e., 0.55 V/cm.). They have stated that at the potential of ECM, the vibration of the DME ceases entirely.

Considering the relatively low intensity of the field in the above study, it appeared to us that the vibration of the mercury drop as a whole may not disturb appreciably the diffusion layer. On the other hand, it was felt that the streaming of the solution by electro-capillaryphoresis may be the cause of the phenomenon. Accordingly, the system was observed through a travelling microscope while illuminating the mercury drop and the solution adjacent to it with

aid of a narrow beam of light from an arc-lamp, condensed with a lens. Some talc powder was added to make any liquid movements present visible.

The system used was the same as investigated by Imai et al.

Solution: 0.001 M Hg^{2+} in 0.1 M KCl.

Drop time: 9.0 Sec. in 0.1 M KCl, open circuit. Alternating field: 0.55 volt cm^{-1} rms.; 50 cycles.

When the alternating field was put on, a well-defined streaming of the liquid took place around the mercury drop. The direction of streaming of solution was found to be the same at all potentials on either side of the ECM. The streaming was from the apex of the drop towards the neck. At the ECM there was no streaming.

We also observed that when a small amount of a surface-active substance like mannoxol OT is added, the streaming of the solution stops altogether at all potentials up to the desorption potential of the surfactant (-1.32 V for mannoxol OT) beyond which the streaming starts again.

In all the above experiments, whenever there was no streaming, normal diffusion currents were obtained. Under similar experimental conditions electro-capillaryphoretic movements were also observed in the case of the hanging mercury drop (the drop adhering to a gold-plated platinum electrode). These movements, however, are not reproducible as the surface appeared to be easily contaminated by surface active impurities.

Differences in interfacial tension appear to be the cause of the well-defined streaming of the solution. The direction of streaming indicates a higher interfacial tension at the neck than at other regions.

It appears to us that the effect of the applied alternating voltage is heterogeneous over the surface of the drop. One of the causes might be that the field would be relatively small at the neck of the drop due to the shielding effect of the capillary and large at the portions which are nearer to the electrodes. As a consequence, different regions are polarised to different degrees. Hence due to electro-capillary effect, the different regions of the mercury drop acquire different interfacial tensions, the neck having a higher interfacial tension than the other regions. Consequently, mercury at the interface flows from the region of low interfacial tension to that of high interfacial tension, i.e., from

apex to the neck; the solution adjacent to the mercury surface flowing along with it thus causing a streaming of the solution.

Thus it appears electro-capillaryphoresis is responsible for the increased limiting current.

In view of the above findings, it is suggested that any mechanical vibratory movement of the drop that may exist may be mainly caused as a result of changes in interfacial tension and may have a subsidiary importance as far as polarographic behaviour is concerned.

Some other related aspects of the phenomenon will be dealt with in a later communication.

One of us (R. N.) thanks the Council of Scientific and Industrial Research for the grant of a Research Fellowship.

Central Electrochemical Research Institute,
Karaikudi (India),
September 7, 1960.

R. NARAYAN,
V. K. VENKATESAN,
K. S. G. DOSS.

1. Imai, H., Inouye, S. and Chaki, S., *Bull. Chem. Soc., Japan*, 1958, **31**, 767; 1959, **32**, 994.
2. Doss, K. S. G., *J. Sci. and Ind. Res.*, 1946, **5B**, 58.
3. Frumkin, A. N., *J. Electrochem. Soc.*, 1960, **107**, 463.

A HEAT-RESISTANT AND ACID-TOLERANT YEAST FROM *PASSIFLORA EDULIS* SIMS.

IN thermal processing or spin-pasteurization of canned fruit products,¹ it is customary to isolate the most heat-resistant organism from the product under test and determine its thermal characteristics with special reference to its thermal-death-time data. During our studies on the isolation of spoilage organisms in passion fruit juice (*Passiflora edulis* Sims.),^{2,3} a survey of literature revealed little published information on the subject. The present report briefly covers this aspect.

Fresh passion fruit juice has, on an average,⁴ 17.3°Brix, 3.4% acidity (as anhydrous citric acid), 2.83 pH, 3.2% reducing sugars, 4.6% non-reducing sugar (sucrose), 10.0% total sugars, 2.4% starch, 0.8% protein, 0.46% mineral matter and 34.6 mg./100 g. of ascorbic acid. Despite its low pH, the fresh juice, when exposed to atmosphere at room temperature, usually got spoiled within 48-72 hours. A microscopic examination of the spoiled juice showed the presence of only yeast cells. The causal organisms were isolated by the

2. Pruthi, J. S., *Ph.D. Thesis*, The Panjab University, Solan, India.
3. — *et al.*, *J. Sci. Industr. Res.*, 1958, **17C** (8), 129.
4. — and Lal, G., *J. Sci. Food and Agri.* (U.K.), 1959, **10** (3), 188.
5. Lodder, J., *The Yeasts—A Taxonomic Study*, Interscience Publishers, New York, 1952.
6. Wickerham, L. J., "Taxonomy of yeasts," *Tech. Bull. No. 1029*, U.S. Dept. Agri., Washington, D.C.
7. Rao, B. A. S. and Pruthi, J. S., *Nature*, 1960, **187** (4734), 342.
8. Lewis, Y. S., *et al.*, *Sci. and Cult.*, 1955, **21**, 220.
9. Pruthi, J. S., *J. Sci. Industr. Res.*, 1958, **17B** (6), 238.

STUDIES ON BOMBAY RATS:

A Note on the Probable Resistance of
B. bengalensis to Plague

The following kinds of rats have been collected from the town of Bombay by different workers.^{1,2}

Rattus rattus, *Rattus norvegicus*, *Bandicota bengalensis*, *Bandicota indica* and *Mus musculus*. On an average 809,744 rats per annum have been brought to this Institute for the last 22 years. In these collections *Rattus rattus* and *Bandicota bengalensis* are the predominating rats. Figure 1 gives the percentage of these two rats in the collections during 1938 to 1960 (June).




FIG. 1. Showing the percentage of the two rats brought to Haffkine Institute from 1937 to July 1960.

R—*Rattus rattus*; B—*Bandicota bengalensis*; A—Human and rat plague cases observed.

From Fig. 1 it will be noticed that the percentage of *B. bengalensis* shows a distinct increase over *R. rattus* and that there is a periodicity noticed in the peak periods. It is interesting to note that whenever the field rats start increasing, the numbers of house rats are going down. This phenomenon may be attributable to the ferocity and size of the field rat, the urbanisation of the town of Bombay and lastly the resistance of *Rattus rattus* and the

susceptibility of *B. bengalensis* to plague.³ There are two known periods marked in Fig. 1 when human and rat plague was in vogue in Bombay. These periods being from 1939 to 1942 and from 1945 to 1951.

TABLE I

Showing the percentage mortality and the average day of death in *B. bengalensis* when submitted to a challenge dose of 195/P.S. *P. pestis* at 5,000 organisms in the laboratory

Species treated	Kamatipura		Mahim	
	% Mortality	Average day of death	% Mortality	Average day of death
<i>R. rattus</i>	17	12.2	27	5.2
<i>Bandicota bengalensis</i>	80	10.1	92	7.1
Controls (Mouse)	100	5.6	100	5.4

Recently *B. bengalensis* were collected in Bombay from the most crowded ward, i.e., Kamatipura and also from an erstwhile suburb of Bombay, i.e., Mahim. Table I gives the percentage mortality and the average day of death as evinced by this rat from the two wards to a standard challenge dose of *P. pestis*. The results indicate the difference in the percentage mortality, confirming the views expressed¹ earlier about the geographical distribution.

The *B. bengalensis* from Kamatipura were later submitted to a standard challenge dose of 10^{-2} , 10^{-3} , 10^{-4} , 10^{-5} dilution of *P. pestis* (195 P-S/in 0.2 ml.). These results indicate that as compared to laboratory mice *B. bengalensis* is 45% and *R. rattus* 96% resistant to plague.

Sera collected aseptically prior to infection from these rats were tested for anti-body to *P. pestis* and *P. pseudotuberculosis* antigens by the agar-gel double diffusion Oudin reaction.^{4,5} It was noticed that the sera from the more resistant rats in Kamatipura displayed anti-body to *P. pestis* and *P. pseudotuberculosis* somatic antigens which are common to both species but not to the specific capsular antigen of *P. pestis*. These observations confirm the experiments of Lawton.^{6,7}

The resistance of *Rattus rattus* and the susceptibility of *B. bengalensis* in Bombay and a number of towns in India to *P. pestis* infection is well known. It is of interest to see that while *B. bengalensis* is replacing *R. rattus* in Bombay it is also in the process of acquiring resistance. Details of these results and serological studies are being published elsewhere.

The significance of these studies are attenuated by the recent positive plague rat fall in Birbhum⁶ district in Bengal where the rat was *B. bengalensis* and the reports of human plague⁹ in the Hosur area of Madras State.

The author is extremely obliged to the Director, Haffkine Institute, for the facilities and to Miss K. K. Patel, Messrs. A. K. Joshi, N. E. Vad, G. C. Chaturvedi and M. K. Habbu for the valuable assistance.


Haffkine Institute,
Bombay-12, August 31, 1960.

P. J. DEORAS.

1. Plague Research Commission, *J. Hyg. Camb.*, **12**, *Plague Suppl.*, 1912, **1**, 229.
2. Deoras, P. J. and Gokhale, M. S., *J. Bom. Nat. Hist. Soc.*, 1958, **55**, 1.
3. *Annual Reports of the Haffkine Institute*, Bombay, 1954, 14; 1955, 16; 1956, 15, 1957, 11.
4. Oudin, J., *Methods in Medical Research*, 1952, **5**, 335.
5. Bhagwan, N. V., Chen, T. H. and Meyer, K. F., *Proc. Soc. Expt. Biol. and Med.*, 1956, **91**, 355.
6. Lawton, W. D., et al., *J. Immunology*, 1960, **84**, 475.
7. Boye, *Bull. Off. inst. Hyg. publ.*, 1932, **24**, 1610.
8. Personal Communication, 1960, Directorate of Health, West Bengal.
9. *Ibid.*, Madras.

SOME REACTIONS OF PHENOLS^[1]

p-CRESOL derivatives are known to undergo the Zincke and Suhl reaction with carbon tetrachloride giving cyclohexadienone derivatives.^{1,2} In the present investigation *p*-cresol has been reacted with carbon tetrabromide in the presence of anhydrous aluminium bromide to yield 4-methyl-4-tribromomethyl-2, 5-cyclohexadienone (I) of m.p. 146°-47° C. (Calc. for $C_8H_7Br_3O$: Br, 66.8; Found: 66.7). The 2, 4-dinitrophenyl hydrazone of (I) gave m.p. 167° C. (Calc. for $C_{14}H_{11}Br_3N_4O_4$: N, 10.4; Found: 10.7). In a similar manner 2, 6-dibromo-*p*-cresol and 3, 4-xyleneol yielded with carbon tetrabromide the corresponding cyclohexadienone derivatives (II) and (III) having m.p. 99°-100° C. (Calc. for $C_8H_5Br_3O$: Br, 77.4; Found: 77.0) and 124°-25° C. (Calc. for $C_9H_7Br_3O$: Br, 64.3; Found: 64.7) respectively. They were characterised by the preparation of crystalline derivatives with 2, 4-dinitrophenylhydrazine.



I: $R_1 = R_2 = R_3 = H$

II: $R_1 = R_2 = Br$; $R_3 = H$

III: $R_1 = R_2 = H$; $R_3 = Me$



(IV)

The compound (I) on a Grignard reaction gave an alcohol (IV), m.p. 177°-78° C. (Calc. for $C_{14}H_{13}Br_3O$: Br, 54.9; Found: 54.6). The latter on treatment with formic acid gave two compounds identified as 4-methyldiphenyl, m.p. 45°-46° C. and 2-methyl-4-phenyl-benzoic acid, m.p. 169°-70° C. Further reactions of the above compounds are being investigated.

p-Ethylphenol and 2, 4-xyleneol were found to react similarly with carbon tetrachloride giving oils from which the 2, 4-dinitrohydrazones prepared had m.p. 169°-70° C. (Calc. for $C_{15}H_{13}N_4O_4Cl_3$: N, 13.3; Found: 13.5) and 166°-67° C. (Calc. for $C_{15}H_{13}N_4O_4Cl_3$: N, 13.3; Found: 13.6) respectively.

A detailed report of the work will be published in due course.

Bombay-1,
Institute of Science,
October 14, 1960.

J. R. MERCHANT.
V. B. DESAI.

1. Newman, M. S. and Pinkus, A. G., *J. Org. Chem.*, 1954, **19**, 978.
2. Newman, M. S., Eberwein John and Wood, L. L. Jr., *J. Am. Chem. Soc.*, 1959, **81**, 6454.

INFLUENCE OF EARTHQUAKE SHOCKS ON THE ASKANIA GRAVIMETER SPRING

THE study of continuous variation in the Gravity value with time has engaged the attention of the authors for a period of over two months (June 28th to September 6th). The Gravimeter is the G_{11} type of the Askani Company and the Gravimeter station is located in the basement of the Geology Department of the Osmania University in Hyderabad (Lat. 17° 26' and Long. 78° 27'). Care has been taken to see that no disturbance is created within the vicinity of the Gravimeter station. During the month of August, disturbances in the general run of the variation curve have been noticed. On the 4th of August at about 1 P.M. a marked drift occurred in the variation curve. For a month prior to this, however, the instrument itself did not show any such or similar drift. In order to explain this sudden change in the behaviour, the authors thought it desirable

FIG.
record
42 sec.

4-8
15-8
23-8
27-8

examine the Seismic records obtained by the Nizamiah Observatory, which is situated about 3 miles West of the Gravimeter station. It was revealed that an Earthquake shock had been recorded at 13 hrs. 17' 42" on the 4th of August, both on the North-South and East-West components. These two records (Gravity and Seismic) stand very well correlated as may be seen from Figs. 1 and 2. Exactly at the same




FIG. 1. The dotted line is the Running Mean Line, and vertical line with dashes, gives the time of the Earthquake shock.



FIG. 2. The North-South component of the Seismic record. Arrow indicates the first shock, at 13 h. 17 m. 42 sec.

time, when the Earthquake shock was recorded by the Seismograph, the Gravity variation curve also showed a marked drift. A detailed examination of the Seismic records showed that the North-South component of the Seismic record showed a more intense disturbance than the East-West component. This shock had a pronounced effect on the Gravimeter spring, which was itself oriented in the North-South direction at the base station, and consequently on the gravity curve. Disturbances, similar to those on the 4th, in the Gravity variation curve and corresponding Seismic records in the North-South component of the Seismogram of the Nizamiah Observatory were noticed on other days of August 1960. The details are given in Table I.

It is possible to conclude from this data that Earthquake shocks felt within the vicinity of the Gravity station have a considerable effect on the continuous Time-Gravity Variation curve. It suggests that the Gravimeter is acting as an Accelerometer. A closer examination suggests the following points:—

1. The direction of the Earthquake shock, with respect to the orientation of the Gravimeter spring, is a factor of considerable importance.

2. The distance of the epicenter from the Gravimeter base station and also the magnitude of the shock are factors that have a marked effect on the behaviour of the spring and hence on the resulting curve.

The authors desire to express their grateful thanks to Professor Masami Hayakawa for his inspiring guidance and many valuable suggestions throughout this investigation.

Geology Department,
Osmania University,
Hyderabad-7 (India),
November 16, 1960.

S. BALAKRISHNA,
P. V. JOHNSON.

TABLE I

Date	Time	Distance of Epicenter	Magnitude	Direction from the Gravimeter station	Effect on the spring
		(km.)			
4-8-1960	13 hrs. 17' 42"	9191	Moderate	Southwards	Marked
15-8-1960	20 hrs. 15' 40"	1765	Slight	N. N-East	do.
23-8-1960	14 hrs. 32' 47"	2420	do.	Southwards	Slight
27-8-1960	21 hrs. 0' 31"	~1200	do.	Northwards	Noticeable

**ON THE EFFECTS OF DIFFERENT
CONCENTRATIONS OF ATMOSPHERIC
OXYGEN ON CELL-DIFFERENTIATION
IN THE REGENERATE OF THE SNAIL,
HELIX ASPERSA MULL.**

MASS (1939) has shown, that the air-breathing snails, when mechanically stimulated, increase their metabolism considerably and, therefore, an increased need for oxygen can be expected. This fact has been confirmed by him in his experiments with *Helix pomatia*.

The object of this investigation was to find out the effect of different concentration of atmospheric oxygen on cell-differentiation during regeneration. At the normal atmospheric condition, the epithelial cells of the regenerate are columnar and the height is 15.6μ . It has been also found that the muscles are in the form of fine strands and are randomly arranged. At the same period, the differentiation of the connective tissue is well advanced. The glands of different types have developed in the regenerate. The free blood-cells are found in numbers but it appears to be migrating from the area into the blood spaces.

below the epithelium have not differentiated so much, as in the normal. The muscle fibres have grown but they are poorly orientated. They are thinner than the normal regenerate.

When different series of animals are treated with 11% O_2 , it is found that at a corresponding stage, the epithelial cells are columnar and closely arranged. Their height has significantly increased to 15.12μ . The developing muscles are not properly orientated. The differentiation of the connective tissue cells is less than the normal regenerate. However, there are masses of blastema and amœbocytes immediately below the epithelium. The hæmocœlic spaces with numerous blood-cells are more than the normal regenerate. The glands have developed and are almost normal.

When animals, after the injury, are kept in 27% O_2 , the epithelium is very similar in shape to that of the normal regenerate but the cells are about 13.3μ in height. A low value of the height of cells, may mean that the epithelium was delayed in the process of growth during that period in high oxygen concentration. The muscle fibres are well developed and have grown

TABLE I
Cell-differentiation in the regenerate of the snail

Atmospheric condition	Epithelium		Connective tissue	Muscles	Glands		
	Shape	Size in Micron			Mucous	Protein	Calcium
1 Normal	Columnar	15.6	Differentiation well advanced	Fine strands randomly arranged	Present	Present	Present
2 With 7% O_2 concentration	Cubical	12.6	Abnormal with oedematous appearance	Grown poorly orientated	22-50% less in number	Absent	Absent
3 With 11% O_2 concentration	Columnar	15.12	Differentiation less pronounced than normal	Developing muscles poorly orientated	100% normal	100% normal	Absent
4 With 27% O_2 concentration	Columnar	13.3	Well differentiated	Muscles grown almost to the normal	Present as normal	Present as normal	Present as normal

It has been observed, that in concentration of oxygen less than atmospheric, the shape and size of the epithelial cells differ from the normal. At 7% oxygen concentration, the cells are cubical and their height is about 12.6μ . The difference in the growth processes of the sub-epithelial tissue is quite marked as compared with the animals grown in normal atmospheric conditions. The important point to be made out here, is that the tissue which is unable to differentiate because of low oxygen tension, may be stimulated to do so by increasing the oxygen concentration. The formative cells

almost to the base of the epithelium, a condition not usually found in normal regenerate. The number and the size of the hæmocœlic spaces have reduced. Different types of glands have been observed in the repaired region.

Barth (1938) has shown that oxygen can be the determining factor in the rate of regeneration in *Tubularia* stems. Tyler (1933) concluded from his work on sea-urchin embryos that differentiation requires the expenditure of metabolically released energy in addition to that required for maintenance. In the present experiment it was observed that the differentia-

No. 1
Dec. 1
tion o
very l
of the
availa
quality
phosis
differ
fore, l
regene
Gold
bitors
play a
in the
the in
in the
A lo
cells, r
during
Anders
of oxy
gress o
playing
regene
cluded
amount
almost
The
Univers
and en
Depart
C.M. C
Darbha

1. An
2. Bar
3. —
4. Gol
5. Ma
6. Ne
7. Tyl

AM
ADU
ANTH
(DE
The fre
Anthren
wool, v
cending
graphy
Adults
starved
mentary
with do
extrane
face of

tion of tissue in low oxygen concentration was very little. The retardation in redifferentiation of the formative cells may be due to inadequate availability of energy. Oxygen affects the quality of the regenerate as in insect metamorphosis (Needham, 1942). It therefore affects differentiation as well as proliferation. Therefore, low oxygen tension was found to retard regenerative processes generally.

Goldin (1942) has shown that certain inhibitors produced by the *Tubularia* stem also play a role during regeneration. An increase in the concentration of oxygen will antagonize the inhibitor allowing more tissue to partake in the process of regeneration.

A low value of the height of the epithelial cells, means that regenerate was delayed even during the period in high oxygen tension. Anderson (1956) has shown that the availability of oxygen acts as a limiting factor in the progress of later process in morphogenesis without playing any necessary part in the initiation of regeneration in *Tubifex*. Barth (1944) has concluded from his work on *Tubularia* that the amount of regeneration at low oxygen was almost equal to a delay at high oxygen.

The author is grateful to Professor A. Graham, University of Reading, England, for guidance and encouragement in this work.

Department of Zoology, B. P. PANDE,
C.M. College, Bihar University,
Darbhanga, Bihar, September 27, 1960.

1. Anderson, J. C., *Biol. Bull.*, 1956, **111**, 179-89.
2. Barth, L. G., *Ibid.*, 1938, **74**, 155-77.
3. —, *Physiol. Zool.*, 1944, **17**, 355.
4. Goldin, A., *Biol. Bull.*, 1942, **82**, 340-46.
5. Maas, J. A., *Zschr. vergl. Physiol.*, 1939, **26**, 605-10.
6. Needham, J., *Biochemistry and Morphogenesis*, Cambridge, 1942.
7. Tyler, A., *Pubbl. Staz. zool. Napoli*, 1933, **13**, 155-61.

AMINO-ACID CONSTITUENTS OF ADULTS OF THE CARPET BEETLE, *ANTHRENUS VORAX* WATERHOUSE (DERMESTIDAE: COLEOPTERA)

The free amino-acid constituents of adults of *Anthrenus vorax* Waterhouse, a serious pest of wool, were qualitatively studied by the descending technique of paper partition chromatography of Consden, Gordon and Martin.¹ Adults of *A. vorax* of both the sexes were starved for 24 hours in order to clear the alimentary canal. They were thoroughly rinsed with double distilled (glass) water to remove extraneous material adhering to the body surface of the insect and were ground with

anhydrous sodium sulphate and 95% ethanol. The homogenates were centrifuged three times at 3,000 r.p.m., each time for about 10 minutes after which supernatants were reduced to 0.2 ml.

As usual the material and standards were spotted on Whatman filter-paper No. 1 (18.25" × 22.5"). Different combinations of *n*-butanol, acetic acid and water were tried and the best separation of free amino-acids was secured by using this solvent mixture in the ratio of 4:1:5² (v/v/v). The chromatograms were run four times and after each run the papers were dried for 6 to 8 hours before running again. After the last run, the chromatograms were allowed to dry overnight and then sprayed with 0.1% ethanolic ninhydrin solution. The chromatograms were dried for an




FIG. 1. Chromatogram of *Anthrenus vorax*.

hour and then kept in dark for 24 hours after which they were kept in an electric oven at 85°-90° C. for ten minutes to develop the spots. The spots were identified by comparing their R_f values with the R_f values of the standards

which were run side by side in the same chromatographic chamber. The entire investigation was carried out at a temperature of $27 \pm 1^\circ \text{C}$, except where mentioned otherwise.

At least the following eleven free amino-acids (Fig. 1): lysine, serine, glycine, glutamic acid, alanine, proline, tyrosine, tryptophan, valine, methionine and isoleucine/leucine were detected by the present technique as the body constituents of the adult carpet beetle of both sexes. Qualitatively there did not appear to be any difference in the constituents of amino-acids in male and female adults.

The writer is thankful to Dr. E. S. Narayanan for his keen interest in this work.

Division of Entomology, SNEHAMOY CHATTERJI.
Indian Agricultural PRAKASH SARUP.

Research Institute,
New Delhi-12, August 11, 1960.

1. Conden, R., Gordon, A. H. and Martin, A. J. P., *Biochem. Jour.*, 1944, **38**, 224.
2. Slotta, K. H. and Primosigh, J., *Nature*, 1951, **168**, 690.

UTILISATION OF THE TECHNIQUE OF VEGETATIVE PROPAGATION IN RICE BREEDING

It is generally observed that under Cuttack conditions, hybrid seeds of artificial crosses lose their viability much earlier than normal seeds, although stored under similar conditions. In a hybridization programme, involving mass production of crossed seeds, hybridization has to be kept going throughout the whole year. Crossed seeds collected during July-October lose their viability by the end of February.

An obvious method of overcoming this difficulty is to grow the hybrid seeds collected during July-October in the second crop season (January to April). The F_1 plants grown during this season, however, show high seed sterility; but the stubbles from these plants can be carried over and grown in the next Kharif season (July to October). The method is also useful in getting enough F_1 seeds for growing large F_2 generation particularly where high sterility occurs.

In research stations, the need is often felt for some F_1 hybrid material either for back crossing to one of the parents, or for verifying some F_2 observations that may have been taken in earlier years. This becomes possible by maintaining F_1 's through vegetative propagation from stubbles year after year. Often interspecific crosses in rice are either almost or completely sterile and the only method available to retain the hybrids for any further work is to maintain

them by stubble planting. This is being done at the Central Rice Research Institute, Cuttack, and a large number of F_1 's of various crosses are being maintained.

STUBBLE PLANTING—AN AID IN THE STUDY OF INHERITANCE OF PHOTOPERIOD SENSITIVITY

In a programme aimed at studying the inheritance of photosensitivity, sensitive varieties are crossed with types, either insensitive or low-sensitive to photoperiod. Since most of the sensitive varieties are short-day plants, F_1 plants are grown in the second crop season (January to April) when the days are longer during the period preceding flowering. If the plants flower during this season, they are taken to be insensitive. To get over the practical difficulty of procuring enough F_2 seeds, the F_1 stubbles are grown in the Kharif season (July to October) and the seeds collected from them are grown in the second crop season. The plants selected from the F_2 generation as either insensitive or low-sensitive are stubble-planted in the Kharif season and the seeds obtained therefrom are used for growing the F_3 cultures in the second crop season.

UTILISATION OF HYBRID VIGOUR

There is marked vigour observed in F_1 's of many crosses in rice in respect of height, number of ear-bearing tillers and yield. One method of taking advantage of the hybrid vigour might be to multiply the F_1 hybrids through stubbles. The best method of propagation is under study and this has been referred to already by the senior author elsewhere.

Central Rice Research Inst., R. H. RICHHARIA.
Cuttack-4, B. MISRO.
August 27, 1960.

1. Richharia, R. H., *Rice News Teller*, 1960, **8**(1), 8.
- 2 —, *Sci. and Cult.*, 1960 (in press).

EFFECT OF CERTAIN ORGANIC INSECTICIDES ON THE YIELD OF CROPS (PART II)

In an earlier paper¹ the writer has described the effect of the application of some synthetic insecticides on the growth and yield of certain crops. In those trials insecticidal treatment was given at more frequent intervals than usual and in the absence of insect infestations. The emphasis was therefore on the reactions, if any, of the plant-body to the insecticidal treatment and not on the effect of such treatment on insect populations affecting the concerned crops.

TABLE I
Results of phytotoxicity trials (Series II); Frequency: weekly

Trial No.	Crop	Area sq. ft.	Insecticide	Number of treatments	No. of plants	Results: yield per plant
1	Potato	27	E, 2 c.c./gal.	9	Tr. 31	2 tubers 27.7 gm.
		C.(30)			C. 28	3.8 " 57 "
2	do.	49	M, 6 "	10	Tr. 27	2.4 " 69 "
		C.(60)			C. 33	2.7 " 78 "
3	Radish	42	F, 3 "	5	Tr. 16	3.3 oz. (with leaves) 1.7 oz. (without leaves)
					C. 23	1.85 " 0.9 "
4	Garden beans	36	F, 3 "	10	Tr. } Not recorded	356 beans 1129 gm.
					C. }	332 " 1088 "
5	Potato	36	G, 5 "	9	Tr. 38	4.9 tubers 37.3 "
					C. 34	2.47 " 23.8 "
6	Radish	15	H, 1 lb./8 gals.	5	Tr. 79	48 gm. (with leaves) 25.3 (without leaves)
			A 10 c.c./gal.		Tr. 64	75.8 " 39 "
			HC 10 "		Tr. 61	73.7 " 37.7 "
			F 3 "		Tr. 46	75 " 36 "
					C. 65	42.3 " 24.6 "
7	Tomato	50	DDT 1 lb./16 gal.	23	Tr. 35	4.8 fruits 94.9 gm.
					C. 24	0.38 " 6.3 "
8	Potato	39	M, 6 c.c./gal.	6	Tr. 28	1.8 tubers 16.6 "
					C. 24	2.6 " 18 "
9	do.	34	*DS 2% "	8	Tr. 26	1.9 " 27.5 "
					C. 14	2 " 16.4 "

Tr. = Treatment; C. = Control.

* Note: Derris Soap was prepared as follows:

Groundnut oil—98 gm.; Derris elliptica powder—5 gm.; Caustic soda—25 gm. (Dissolved in 100 c.c. water); Resin..... a small lump. The oil was heated on a low flame. The other ingredients were added to the warm oil and thoroughly stirred in.

In the present note results of a further set of trials conducted between November 1958 and September 1959 along the same lines are presented. In this series there were 10 trials relating to nine insecticides, viz., Folidol (F) (3 trials), DDT (E) (2 trials), Malathion (M) (2 trials), DDT 50% W.P. (2 trials), Gusathion (G), Mangay oil soap (H), 2% Derris soap* (DS), Aldrin (A) and Heptachlor (HC). The crops sown with were Potato (5 trials), Radish (3 trials), Garden beans, Tomato and groundnut. The first nine trials were laid out in the laboratory compound in small plots varying in size from 15 to 50 sq. ft. and the last trial (on groundnut) in Ramakrishna Krishi Shala (Anekal Taluk) on duplicated 2-guntha sub-plots. There were check plots of approximately the same size in all cases. The crops were allowed to remain till the end, being harvested at the end or at intervals, as required.

In Tables I and II are given the data relating to these trials:—

In many of these trials spraying with insecticides has affected the yield one way or the other. Where there is an improvement in yield over the control it is obviously due to factors

TABLE II
Results of phytotoxicity trials on groundnut (Trial No. 10)

Crop: Groundnut. Sub-plot size: 2 guntha.
Sown: 17-6-1959. Harvested: 14-11-1959.
Insecticides:—(a) Aldrin .. 10 c.c. per gallon.
(b) Heptachlor .. " "
(c) DDT 50% W.P. 1 oz. per gallon.
(d) Folidol .. 2 c.c. per gallon.

Each insecticide was applied in two sub-plots as shown in the following sketch:

A	B	C	D	E
Aldrin	Heptachlor	DDT	Folidol	Control
F	G	H	I	J
Control	Folidol	DDT	Heptachlor	Aldrin

Frequency: Fortnightly.

No. of treatments: 9

Harvest data

Treatment	Average weight of pods harvested (lb.)	Variation from check	By measure (Mysore seers)	Variation from check
Aldrin	52	+ 2	31.8	+ 0.3
Heptachlor	32½	-17½	20.5	-11
DDT 50%	51½	+ 1½	32	+ 0.5
Folidol	56½	+ 6½	35½	+ 4
Control	50	..	31½	..

other than pest control since no infestation was observed in these trials. Similarly where the yield is less than in the control, treatment should be considered as one of the factors responsible for it. In either case it may be permissible to infer that the concerned insecticides are absorbed by the plant-body and influence the yield.

Ekatox and Malathion (2 trials) both on Potato, have given adverse results. Similarly Heptachlor on groundnut used in combination with 3 other insecticides has not given favourable results. Spraying with the other insecticides has resulted in increased yields. The performance of DDT 50% W.P. on Tomato (Trial No. 7) deserves mention here. The treated sub-plot has given 4.8 fruits weighing 95 gm. per plant, while the control has yielded 0.38 fruit weighing 6.2 gm. per plant. This large increase under insecticidal treatment requires confirmation by further tests. Hongay oil Resin soap and Derris soap which are in a category different from the other organic insecticides have given encouraging results.

Grateful acknowledgements are due to Dr. M. Puttarudriah, Government Entomologist, for encouragement and advice, and to Mr. D. P. Ramanna, Laboratory Assistant, for help in field-work.

Division of Entomology, D. SESHAGIRI RAO.
Department of Agriculture,
Bangalore, June 3, 1960.

1. Seshagiri Rao, D., *Curr. Sci.*, 1959, 28 (2), 57.


A HAEMATOXYLIN SQUASH METHOD FOR THE ROOT TIPS OF *DOLICHOS* *LABLAB* LINN.

THE meagre information available on the nuclear cytology of *Dolichos lablab* Linn. is not commensurate with the economic importance of the species.¹⁻⁵ The chromosomes are small and the meristematic cells are often refractory to the application of conventional squash methods. This necessitated a search for a simple technique capable of application to such difficult material.

The limitations of aceto-carmin and aceto-orcein techniques are well known.^{6,7} Tiwary and Shrivastava⁸ devised an aceto-haematoxylin method and claimed that it gave pictures superior to those obtained with aceto-carmin. Melander and Wingstrand⁹ used Gomori's haematoxylin and obtained squashes after treatment with 45% acetic acid. They claimed a high selectivity for the stain and considered it superior to others for difficult material. The limitations of the technique were the variation

from material to material of the time of hydrolysis in NHCl at 60° C. prior to staining, the dependence of the affinity of the cytoplasm on the composition of the fixative used and the necessity often of the use of a counterstain for the cytoplasm.

The technique given below stains all cell organelles in differing shades of blue. The root tips of *D. lablab* fixed in acetic alcohol (1:3) for 1-24 hr. were either stored in 70% alcohol or used immediately after a thorough wash in distilled water. They were hydrolysed in NHCl at 60° C. for 10-12 min., washed in distilled water, mordanted in 4% ferric ammonium sulphate for 5-10 min., washed in distilled water and stained for 15-20 min. in well ripened haematoxylin. They were then washed for 10-15 min. in distilled water and softened with 45% acetic acid at 60° C. To obtain good squashes, the primary roots had to be kept in acetic acid for 8-10 min. while the secondary roots required only 4-5 min. Since acetic acid is the destaining agent, the time of stay in haematoxylin had to be adjusted to the time required for softening in acetic acid.



FIGS. 1-4. Fig. 1. Resting Nucleus, \times ca. 1,700. Fig. 2. Prophase, \times ca. 2,000. Fig. 3. Polar view—Metaphase, \times ca. 2,400. Fig. 4. Polar view—Note the pair of satellites, \times ca. 3,000.

The root tips washed in distilled water were mounted in a drop of 45% acetic acid under coverslip, squashed and the preparations sealed

with p
make
remove
butyl
slip we
in Cam

Figure

nucleob
in Fig.

(Figs. 3
instead

Attentio
chromos

The 1

Phaseob
unpubli

cepa (S

Cytogen

Dept. of

Indian I

Bangalo

1. Karp

2. Kaw

3. Ayya

4. Rau

5. Roy

6. Joha

7. Tandi

8. Tiwar

9. Melan

10. Meena

11. Sen

196

A CASE

At the V
case of
March, 19
planted in
a bulb crop
the crop v
One of
demarcate
the other a
green sect
meter of l
the leaves
clearly. L
bulb crop, s
sprung up
ing shoot a

with paraffin wax for a general survey. To make them permanent, the paraffin seal was removed and the slide transferred to tertiary butyl alcohol. The separated slide and cover-slip were further dehydrated and then mounted in Canada balsam.

Figure 1 shows the resting nucleus with the nucleolus and the early prophase is illustrated in Fig. 2. A polar view of the metaphase plate (Figs. 3 and 4) shows only 22 chromosomes^{1,2,5} instead of the 24 reported by some workers.^{3,4} Attention is invited to a pair of satellited chromosomes¹¹ (indicated by arrows in Fig. 4).

The technique gives equally good results in *Phaseolus radiatus*, *Pisum sativum* (Royan, unpublished), *Cicer arietinum*¹⁰ and *Allium cepa* (Subramanyam, S., unpublished).

Cytogenetics Laboratory, K. M. MARIMUTHU.
Dept. of Biochemistry, M. K. SUBRAMANIAM.
Indian Institute of Science,
Bangalore-12, October 12, 1960.

1. Karpechenko, *Bull. App. Bot.*, 1924-25, **14**, 143.
(Quoted by Ayyangar and Krishnaswamy.)
2. Kawakami, J., *Bot. Mag. Tokyo*, 1930, **44**, 319.
3. Ayyangar, G. N. R. and Krishnaswamy, N., *Curr. Sci.*, 1935, **4**, 739.
4. Rau, N. S., *J. Ind. Bot. Soc.*, 1929, **8**, 201.
5. Roy, P. and Saran, J., *Proc. Ind. Sci. Cong. Abstr.*, 1959, **46**, 350.
6. Johansen, D. A., *Plant Microtechnique*, McGraw Hill Book Co., Inc., New York, 1940.
7. Tandler, C. J., *Stain. Tech.*, 1959, **34**, 234.
8. Tiwary, N. K. and Shrivastava, S., *Sci. and Cult.*, 1950, **16**, 163.
9. Melander, Y. and Wingstrand, K. G., *Stain Tech.*, 1953, **28**, 217.
10. Meenakshi, G. and Subramaniam, M. K., *Curr. Sci.*, 1960, **29**, 438.
11. Sen, N. K., and Marimuthu, K. M., *Caryologia*, 1960, **13**, 411.

A CASE OF A SECTORIAL CHIMERA IN ONION

At the Vegetable Research Station, Solan, a case of *Chimera* in onion was observed in March, 1960. The seedlings had been transplanted in November-December 1959 for raising a bulb crop. The Development and growth of the crop was normal.

One of these plants showed two, clearly demarcated, longitudinal sectors one green and the other albino as seen in the photograph. The green sector was about two-third of the diameter of leaves and albino about one-third. All the leaves of the plant showed the character clearly. Later in April, as is common in this bulb crop, some of the plants, as also the *Chimera*, sprung up early flowering shoots. The flowering shoot and the unopened umbel in the parti-

cular plant showed the same demarcated sectors of albino and green.




FIG. 1. Taken at the time of springing up of the flowering shoot.

During a severe storm the flowering shoot however broke. The bulb has been carefully stored and will be planted for seed production in the end of the year. The breeding behaviour of the plant will be studied from the seeds and the bulb saved in the second year.

Assistant Vegetable Botanist, S. L. JUNEJA.
Himachal Pradesh, Solan, RAM GOPAL.
July 26, 1960.

A NEW SPECIES OF *CERCOSPORA* ON *LEPTADENIA RETICULATA* WIGHT AND ARN.

Leptadenia reticulata, a common climber, was found to be heavily infested with a species of *Cercospora* in the month of October, 1959, near Udaipur City. Since the fungus has never been reported on this host, a brief description of the causal organism and the symptoms which it causes are given in what follows.

The disease starts with the appearance of olivaceous brown to fuliginous patches on the lower surface of the leaves, presence of which is indicated on the other surface by pale yellow

discolouration in later stages. Large number of such spots of irregular size appear on the leaves which often coalesce and cover the major portion of its surface. Spots also appear on green parts of the stem and branches. In advanced stage of the disease, leaves turn yellow, dry up and ultimately fall off.

Mycelial patches are amphigenous, effuse, irregular, first scattered but later coalesce, olivaceous brown in colour, highly variable in diameter, mostly on green parts of stem and lower surface, rarely on upper surface of the leaves.

Conidiophores fasciculate, arising from submerged synnema, pale olivaceous or olivaceous brown, uniform in colour, somewhat irregular in width, longer ones curved, unseptate to one septate at the base, tip bluntly rounded, spore scars small, measuring 12 to 28.8 μ \times 3.6 to 6 μ with an average of 21.6 \times 4.8 μ (Fig. 1A).




FIG. 1. A—Transverse section of leaf showing conidiophores and conidia. B—Conidia.

Conidia subhyaline to very pale olivaceous, thickwalled, one to more septate, cylindro-obclavate, straight to curved, tip subobtusate to obtuse, fragile, measuring 29 to 126 μ \times 3.6 to 5.4 μ with an average of 59 \times 4.4 μ (Fig. 1B).

On green leaves and stem of *Leptadenia reticulata* Wight and Arn. Leg. R. D. Singh, G. C. Bhatnagar.

Type specimen deposited in the Herbarium of C.M.I., Kew, England, and Herb. Crypt. Ind. Orient, I.A.R.I., New Delhi.

Since no report of any *Cercospora* occurring on *Leptadenia reticulata* Wight and Arn. could be traced out from the available literature, the name *Cercospora leptadeniae* has been proposed to recognize it as a new species.

Sincere thanks are due to Dr. J. C. F. Hopkins for confirming the identification and to Shri Samarath Raj, Director of Agriculture, Rajasthan, for facilities.

Plant Pathology Section,
Department of Agriculture,
Rajasthan, Udaipur,
April 7, 1960.

N. PRASAD.
R. D. SINGH.
G. C. BHATNAGAR.

1. Chupp, C. A., *A Monograph of the Fungus Genus Cercospora*, 1953.

PRELIMINARY STUDIES ON THE EFFECT OF GIBBERELIC ACID ON GROWTH OF SPINACH (*SPINACHIA OLERACEA*)

GIBBERELIC ACID (GA) is believed to promote the overall expansion and elongation in a wide variety of plant species.¹⁻³ An attempt was, therefore, made to study the influence of GA on a highly nutritive leaf crop like spinach (*Spinacia oleracea*).

There were six treatments (including control) with three replications of each. Seeds were sown in beds on December 19, and seedlings were thinned to one foot distance (plant to plant and row to row). Seedlings were given three drench sprays of different concentrations of GA (0; 10; 25; 50; 100 and 250 parts per million) at 40, 55 and 70 days after sowing, in randomly selected beds. Control plants were sprayed with distilled water. Height of the main shoot, number of branches and leaves per plant, fresh and dry weight of foliage per plant were recorded 90 days after sowing (Table I).

TABLE I

The effect of different concentrations of GA on the average and ultimate height of plant, number of branches, number of leaves and fresh and dry weight per plant in spinach

GA concentrations in parts per million	Average ultimate height per plant in cm.	Average No. of branches per plant	Average No. of leaves per plant	Average fresh weight per plant in gm.	Average dry weight in gm.
0 (Control)	38.66	2.17	24.66	44.77	12.10
10	90.83	14.66	50.30	378.38	80.18
25	82.83	11.00	42.66	213.06	45.06
50	75.60	11.66	36.33	223.00	50.00
100	70.83	8.43	32.00	101.00	25.76
250	55.20	6.23	28.00	80.00	25.00

C.D. at 5% 13.37 2.47 7.05 19.49 8.36

Table I suggests that different concentrations of GA brought about a marked improvement in the growth of the plants. The maximum average height (90.83 cm.) with 14.66 branches

and material increase in the number of leaves per plant was obtained under 10 p.p.m. (Fig. 1).




FIG. 1. Influence of different concentrations of GA on the growth of spinach. (left to right) 10 p.p.m., 50 p.p.m., 100 p.p.m., 250 p.p.m. and Control.

Treated plants produced a much higher fresh and dry weight per plant, as compared to untreated ones. Significantly higher fresh and dry weights (378.38 and 80.16 gm. respectively) were also achieved under 10 p.p.m.

It is, thus, obvious that the foliar sprays of the lower concentrations of GA (10 p.p.m.) may be conveniently employed for increased production of spinach.

Govt. Agric. College,
Kanpur, July 6, 1960.

O. S. JAUHARI.
R. D. SINGH.
V. S. DIKSHIT.

1. Brian, P. W. and Grove, J. F., *Endeavour*, 1957, 16, 161.
2. Bukovac, M. J. and Wittwer, S. H., *Michigan Agric. expt. Sta. Quart. Bull.*, 1956, 39, 307.
3. Marth, P. C., Audia, W. L. and Mitchell, J. W., *Bot. Gaz.*, 1956, 118, 106.

THE PRESENCE OF INDOLE-LIKE COMPOUNDS IN PARTHENO-CARPIC GUAVA

In a recent communication from this laboratory parthenocarpic development of fruits in guava, *Psidium guajava* L. var. 'Allahabad Round', induced by 'pollen hormone' was reported.¹ It was found that the water extract of pollen grains, when applied to emasculated flowers, was capable of causing parthenocarpic fruits. In order to study the biochemical aspects of the problem, further investigations were made and results are reported here.

Freshly collected pollen grains were extracted, following the method described by Wright,² with slight modifications. The pollens were treated for 24 hours with two changes of peroxide-free ethyl ether at 0° C., using 50 ml. of ether per gram of pollen. The ether extract was separated and evaporated to dryness over a water-bath at 60° C. The residue was mixed in lanolin paste and used for treating emasculated flowers. Synthetic auxins were also applied to the flowers in the same manner. The results obtained are summarized in Table I.

The water as well as ether extracts of pollen were found capable of causing parthenocarpic development of the fruits, and of the synthetic auxins tested, NAA, NOA and 2, 4-D were found effective in causing parthenocarpy.

In order to examine the extract for the presence of any indole compound further work was carried out. The flower parts, viz., pollen grains and ovaries at different stages of growth were selected and the ether extracts prepared as stated earlier. The extract was distilled off at 60° C. and the residue digested at 70° C. for five minutes with small quantities of distilled water. The digested material was kept at low temperatures for a day after which the solidified fatty substance containing chlorophyll was filtered off. The filtrate was extracted repeatedly with ether. The ether extracts thus obtained were bulked and treated with one-sixth the volume of 5% sodium bicarbonate solution and the two layers separated with a separating funnel. This process was repeated 5 times. By this process the acidic substances are brought to the bicarbonate fraction, while the non-acidic substances remained in the ether fraction. The bicarbonate fractions were bulked and acidified to pH 3 with 20% tartaric acid and re-extracted three times with equal volume of ether. Then the ether extracts containing acidic fraction were bulked and dehydrated with anhydrous sodium sulphate and concentrated over a water-bath at 60° C. and used for chromatographic differentiation. In the same manner the non-acidic fraction of the extract was concentrated and used for chromatographic differentiation. Both circular and ascending chromatograms were developed with Butanol-ammonia-water (100:100:8, upper layer) with Whatman No. 1 filter-paper in complete darkness. The indole compounds on the chromatograms were detected by spraying modified Salkowski reagent (10 parts of 5% perchloric acid and 1 part of 0.05 M ferric chloride). Known auxins were used for comparison. Only the acidic fractions were found to contain indole-like compounds and not the

TABLE I
Inducement of parthenocarpy in guava

Treatment	No. of flowers treated	No. of flowers set	Remarks
Water extract of the pollen	60	26	The set fruits were parthenocarpic; the unset flowers dropped off in 12 to 15 days.
Ether extract of the pollen	120	55	The set fruits were parthenocarpic, but a majority of them dropped off in 20 to 25 days.
Naphthalene acetic acid (NAA) 0.5%	30	10	The set fruits were parthenocarpic
Naphthoxy acetic acid (NOA) 0.5%	30	8	do.
2, 4-Dichlorophenoxy acetic acid (2, 4-D) 0.5%	30	6	do.
3-Indole acetic acid (IAA) 0.5%	30	0	There was no set and the flowers dropped off in 12 days.
3-Indole butyric acid (IBA) 0.5%	30	0	do.
Self-pollinated and bagged	60	15	The set fruits were parthenocarpic
Emasculated and bagged	60	0	All the flowers dropped off in 12 days.

non-acidic fractions, as detected on the chromatograms.

By this method traces of an indole-like compound were detected in the pollens collected immediately after the opening of flowers. In the case of ovaries there was no indication of detectable quantity of the chemical at the time of flower opening, but considerable quantity could be detected in the ovaries collected 5 days after selfing; two fractions of the compound with different Rf values were detected on the chromatograms (Fig. 1). But developing



FIG. 1. Chromatographic separation of the ether extracts of guava ovary: G.O.: acidic fraction; N.: non-acidic fraction; K: known compound—IAA; 1 and 2: two fractions of indole-like compound found in the acid fractions of the ether extract.

young fruits, 20 days after selfing, did not contain any detectable quantity of the indole-like compound. When the flower parts of a seeded variety of guava, viz., 'Red-fleshed' was examined for the purpose, in no case the chemical could be detected on paper chromatograms.

The indole-like compound present in the pollen grains, and ovaries of 'Allahabad Round' variety of guava gave a pink colour on chromatograms when treated with modified Salkowski reagent, which is the same as that of known IAA. But the Rf values of the two unknown fractions were 0.93 and 0.64 as against 0.57 for IAA. It is probable that the fractions are closely related in their molecular structure to IAA. But IAA when applied to the flowers was found ineffective in causing parthenocarpic development of the fruits whereas NAA, NOA and 2, 4-D were fairly effective in this respect (Table I). The colour and the Rf values of the latter chemicals on chromatograms were, however, different from those of the unknown fractions as seen in the present studies. The exact identity of these compounds and their relationship to parthenocarpic development of guava remains to be investigated.

Dept. of Agriculture, G. RANGASWAMI.
Annamalai University, T. T. KALIAPERUMAL.
Annamalainagar,
September 5, 1960.

1. Balasubramanian, V. R. and Rangaswami, G., *Curr. Sci.*, 1959, 28, 413.
2. Wright, S. T. C., *J. Hort. Sci.*, 1956, 31, 196.

FA
SORGH

AMONG t
Cholam)
tillering
solitary
large an
other m
condition
in wild
tillering
level, wh
and coul
develop
distinct f
shoots ar
possible
basal bud
nected w
Vellai Ch
The m
straw qua
and a wh
as a simp
non-tiller
were cros
former to
summer s
subsequen
There was
plants we
in running
postings f
together
presented

Counts fo

Non-tillered
Tillered
Total

The obs
indicating
inhibitory
strains sho
ing charact
as it exhib
when indu
wider spac
The tiller
also (Fig. 1
etting goo
new vigor
main shoot.

**FACTORS FOR TILLERING IN
SORGHUM SPP. (S. SUBGLABRESCENS)**

AMONG the millets in *S. subglabrescens* (Vellai Cholam) and certain other species of *Sorghum* tillering is conspicuously absent. However, the solitary earheads borne on the single stalks are large and heavy compared to earheads of the other millets that tiller well. Single-stalked condition in grain sorghum differs from that in wild sorghum. In the latter the absence of tillering is due to absence of buds at the ground-level, whilst in the former the buds are present and could be activated.¹ The basal shoots that develop from the buds at the ground-level as distinct from the main stem and other secondary shoots are termed the tillers.² The factors responsible for suppressing the development of the basal buds were revealed during studies connected with improvement of straw quality of Vellai Cholam through hybridization.

The midrib of sorghum is an index of the straw quality. Dull midrib indicates juicy straw and a white one pithy straw and white behaves as a simple dominant over the dull type.³ Two non-tillering Cholam strains Co. 18 and K 2 were crossed to transfer the juicy stalk of the former to the latter. The cross was effected in summer season 1956 and F 1 and F 2 studied in subsequent years. The F 1 was single-stalked. There was segregation in the F 2 and the tillered plants were carefully levered out and washed in running water to confirm this habit. The postings for tillered and non-tillered plants together with their statistical significance are presented in Table I.

TABLE I

Counts for	Observed	Expected	X ²	Probability
Non-tillered ..	495	504.5625		
Tillered ..	126	116.5775	0.8750	70.3640
Total ..	621	621.0000		

The observed figures agree to a 13:3 ratio indicating that one of the parents carries an inhibitory factor. Past experience with the strains shows that Co. 18 might carry the tillering character as well as the inhibitory factors as it exhibits a greater proclivity for tillering when induced by ratooning, insect damage or wider spacing.

The tillering habit appears to be economical also (Fig. 1). The earheads are bold and grain setting good. The tillers ranging from 1 to 5 grew vigorously and matured along with the main shoot. Apart from the difference in num-

ber of tillers there was no other perceptible variation amongst the tillered plants. The harvest can be completed in one lot as maturity is uniform unlike in the millets cumbu and ragi which also tiller freely.




FIG. 1. Tillering and non-tillering conditions in Vellai Cholam. 1. A clump from F 2 of Co. 18 x K 2 with five tillers. 2. Single-stalked shoot of cholam K 2 (Co. 18 being similar to K 2 for tillering it is excluded from the photograph, vide text for details.)

The valuable guidance rendered by Sri. B. W. X. Ponnayya, Professor of Genetics, Agricultural College and Research Institute, Coimbatore and Sri. S. G. Ayyadurai, Millets and Pulses Specialist, in presenting the paper, is gratefully acknowledged.

Agricultural Res. Station, K. DIVAKARAN.
Kovilpatti, July 26, 1960.

1. Ayyangar, Rangaswami, G. N. and Ponnayya, B. W. X., *Proc. Ind. Acad. Sci.*, 1939, 10 B (4), 237.
2. Goodal, S. V., *Mem. Dept. Agri. in India, Botany*, 1928, Ser. XIX, 247.
3. Hilson, G. R., *Agric. Jour. Ind.*, 1916, 2, 150.

**PROPAGATION OF *ALTHEA ROSEA*,
BENTH. AND HOOK. (HOLLY HOCK)
BY AIR LAYERING WITH THE AID OF
GROWTH REGULATORS**

Althea rosea is an important ornamental shrub, which has so far been propagated by seed. It is difficult to raise plants true to type by seed propagation. Therefore, an attempt was made to propagate the plants by air layering with the aid of IBA, NAA and mixture of both.

Side shoots of 2-3 months old plants were selected for layering purpose. Indole butyric acid and Naphthalene acetic acid at concentrations of 4,000 p.p.m., 3,000 p.p.m. and 2,000 p.p.m. were used individually and in combination of both 1 mg. IBA + 1 mg. NAA with 1 gram of active carbon powder. The hormones were first dissolved in alcohol and mixed with carbon. The mixture was dried and powdered well. About 3 cm. wide bark was peeled off in the form of a ring. Upper cut end of the ring was smeared uniformly with different concentrations of hormones by means of camel-hair brush after moistening the ringed portion with water. Shoots treated were covered with damp soil containing leaf mould and tied firmly with Alkathene.

TABLE I


The effect of IBA, NAA and combination of both on rooting percentage, nature of rooting and average length of longest root in the air layer of *Althea rosea*

Sl. No.	Hormones in p.p.m.	Percentage of rooting	Nature of rooting	Average length of the longest in cm.
1	IBA			
(a)	4,000	100	S	5.4
(b)	3,000	100	P	9.7
(c)	2,000	100	P	7.5
2	NAA			
(a)	4,000	100	S	10.8
(b)	3,000	100	P	10.06
(c)	2,000	100	P	10.4
3	IBA+NAA + 1 gram A.C.	100	P	10.5
4	Control	30	N	..

Notes: P=Profuse, S=Sparse, N=Negligible.
A.C.=Active Carbon.

There were twelve shoots under each treatment. Gootes were examined for rooting at six days interval. Twenty-two days after actual treatment, gootees were cut-off from mother plant for final observations.

All the treatments with regulators proved advantageous and cent per cent success was achieved in gootees (see Fig. 1 and Table I).



FIGS. 1 a-b. Fig 1 (a). A gooteed plant of *Althea rosea*. Fig. 1. (b) Showing rooted twigs under each treatment with different kinds of hormones.

The lower concentrations of both hormones were found favourable in inducing roots. The higher concentrations were less effective. The lower concentrations induce roots as early as 14 days from the date of treatment.

Author is grateful to Shri B. Venkoba Rao Principal, for his interest and providing facilities.

Division of Botany,
College of Agriculture, Hebbal,
Bangalore, August 16, 1960.

D. S. LINGARAO

INSECT

THE s
is one
distribu
India,
sometim
harvest
control
endrin,
at Anah
Subsequ
lindane
conclusi
BHC a
seemed
compara
(all 5%
at a ra
gradient
was stu
(Patna d
in Table

Effective

Treatme

Infecticide
BHC
BHC
Chlordan
Chlordan
Parathion
Parathion
Aldrin
Aldrin
Control

Aldrin
festation
acre was
treatment
also high
than yield
1 pound
of infeste
of the tre
were also
most ineff

No phy
plots trea
these plots
taste or fl

**FIELD TRIALS WITH SOME
INSECTICIDES FOR THE CONTROL OF
SWEET POTATO WEEVIL**

THE sweet potato weevil *Cylas formicarius* F. is one of the most destructive and widely distributed insect pests of sweet potatoes in India. Damage caused by it in Bihar alone sometimes extends up to 90% of the late-harvested crops. Recent experiments for the control of this pest with lindane, aldrin, dieldrin, endrin, DDT, parathion and calcium arsenate at Anakapalle (Andhra) yielded erratic results.¹ Subsequent trials with aldrin, dieldrin and lindane in the Madras State also proved inconclusive.² Only in Mysore State, dusting with BHC and spraying with parathion (Folidol) seemed to have afforded partial relief.³ The comparative efficacy of BHC, aldrin, chlordane (all 5% dusts), and parathion (20% wettable) at a rate of one and two pounds (active ingredients) per acre, against *Cylas formicarius* F., was studied in a randomized field trial at Patna during 1957-58. The results are given in Table I.

TABLE I
Effectiveness of insecticides applied for the control of *Cylas formicarius* F.

Treatments	Rate of application per acre	Weight of tubers per plot in oz.	Percentage infestation	Effectiveness*
Insecticide (pounds)	Healthy	Infested	%	%
BHC .. 1	230	23	9	74.4
BHC .. 2	223	20	8	77.7
Chlordane .. 1	298	25	8	72.2
Chlordane .. 2	194	29	13	67.7
Parathion .. 1	223	69	24	23.3
Parathion .. 2	250	51	17	32.2
Aldrin .. 1	294	22	7	75.5
Aldrin .. 2	309	12	3	86.6
Control	219	90	29	..

*By Abbott's formula.

Aldrin gave the best results. Average infestation in plots having 2 pounds of aldrin per acre was only 3% and the effectiveness of the treatment was rated at 86.6%. The yield was also highest—411 ounces per plot (82% higher than yield in control plots). In the plots having 1 pound aldrin per acre, the average percentage of infested tubers was 7 and the effectiveness of the treatment was 75.5%. BHC and chlordane were also effective, but parathion proved to be most ineffective of all the insecticides used.

No phytotoxic effects were evident in the plots treated with aldrin. The tubers from these plots did not exhibit any deterioration in taste or flavour.

Grateful thanks are due to Dr. Pushkarnath, Director, for his kind interest in the work.

Central Potato Research Institute, Simla,
K. K. NIRULA.
K. N. CHHIBBER.
Potato Research Station, P.O. Sahay Nagar,
Patna (Bihar), August 29, 1960.

1. *Annual Progress Report of the Tuber Crop Scheme in Andhra State, Anakapalle, for 1954-55.*
2. *Annual Progress Report of the Tuber Crop Scheme for 1954-55 in Madras State.*
3. *Final Report of the Scheme for Research on Tuber Crops for the Period 1st July 1951 to 31st March 1955.*

**SHELL DISEASE IN CRASSOSTREA
GRYPHOIDES (SCHLOTHEIM)**

SHELL diseases have been prevalent in British, Dutch and French oysters. Cole and Waugh (1956) find that in case of British oysters, the disease commences with the dendritic white markings in the shell, usually in both valves. These markings are probably the results of infection by a fungus. The disease occurring in Dutch and French oysters is commonly referred to as 'maladie du pied', and here, it is in the form of greenish-brown patches and warts, leading to the serious loss of condition. When the disease reaches the muscle attachment, there is interference with the closing mechanism, resulting in death.

The occurrence of a shell disease does not appear to have been recorded in case of the back-water oyster, *C. gryphoides*. However, during the collections of *C. gryphoides* from a cultivated farm near Bombay for the study of some aspects of its biology, seven specimens were noted with shell disease. The interior of the shell of one oyster was strewn with greenish-brown patches and the muscle scar appeared disfigured (Fig. 1). The disease of this oyster, therefore, appears to resemble, 'maladie du pied', diagnosed by Giard (1894), Hornell (1910), Ranson (1936) and Cole (1950) in French and Dutch oysters.

In the case of five other oysters, a dark-green patch on the interior of the shell was noticed, but there was no marked disfiguration of the muscle scar or the interior of the shell as mentioned above (Fig. 2). It is likely that this may perhaps be the beginning of the oyster disease. In the case of the remaining specimen of oyster, the muscle scar was found blistered and the animal was almost detached from its shell. However, the interior of the shell was not disfigured by greenish-brown patches as above (Fig. 3).




FIG. 1

In all these specimens, the meat was watery, pale and brown in colour, giving the oyster unhealthy appearance.

Korringa's investigations have shown that shell disease spreads rapidly in hot summers. However, in case of *C. gryphoides* out of nearly three thousand specimens examined during a period of two years only seven were found affected, indicating thereby the absence of the spread of shell disease.

Department of Zoology,
Institute of Science,
Bombay-1, September 22, 1960.

V. S. DURVE.
D. V. BAL.

1. Cole, H. A., *Nature*, 1950, **166**, 19.
2. — and Waugh, G. D., *Ibid.*, 1956, **178**, 422.
3. Giard, A., *C.R. Soc. Biol.*, 1894, **46**, 491.
4. Hornell, J., *Madras Fish. Bull.*, 1910, No. 4, 1-31.
5. Korringa, P., *Cons. prem. Int. pour l'explor. de la mer.*, Edinburg, 1949.
6. Ranson, G., *C.R. Soc. Biol.*, 1936, **121**, 540.

* Not referred to in original.

OCCURRENCE OF *THOREA RAMOSISSIMA* BORY. IN INDIA

The genus *Thorea* Bory (Thoreaceae, Nemalionales, Rhodophyta) comprises about 6 species all of them occurring in fast flowing freshwater streams. The type of genus is *Thorea ramosissima* Bory. The genus is known from Europe, America, Japan and some of the larger islands in the Indian Ocean (Fritsch, 1945; Kylin, 1956). This genus has not been reported from India so far. From India only some other species of freshwater genera of the Rhodophyceae namely *Batrachospermum* and *Compsopogon* have been recorded.

The author while making algal collection, in a swiftly running stream Narkatia near Bareilly, in the last week of January, 1955, encountered the alga for the first time (Fig. 1). During subsequent years, i.e., 1956, 1957 and 1958, regular collections from the same spot have been made. The plants show fairly luxuriant growth and attain a length of two to three feet. *Compsopogon caeruleus*




FIG. 2

Mont. and a species of *Batrachospermum* were also collected from the same rivulet. During regular collections, however, it has been observed that *Batrachospermum* is the first to appear in



FIG. 3



FIG. 1. Whole mount of the alga showing the thallus and the threads covering it, $\times 30$.

December, it starts degenerating by February, whereas *Thorea ramosissima* appears last and continues a luxuriant growth even up to early March.

The author is thankful to Dr. P. Bourrelly, Director, Museum National D' Histoire Naturelle, Laboratoire De Cryptogamie, Paris, for confirming the plant as *Thorea ramosissima* Bory, and to Dr. Bahadur Singh for going through the manuscript.

University Dept. of Botany, DALBIR SINGH,
Jaswant College,
Jodhpur, August 31, 1960.

1. Fritsch, F. E., *The Structure and Reproduction of the Algae*, 1945, Vol. II, London.
2. Kylin, H., *Die Gattungen der Rhodophyceen, Glaurops, Lund.*, 1956, p. 101.

Report
Edito
Societ
Road
63 sh
The
Physic
been r
quarte
part of
all coll
possess
forward
A welco
the Soc
bound v
separate
wider u
particul
objects
general
the ann
this rev
content
the tem
briefly i
them.

The fi
thin film
the auth
of a sur
dimension
the stru
as aggre
orientati
extents
millimic
electrom
a refract
an extinc
discussio
importan
The so
solid st
(pp. 66-
photomet
and the
analysis
(pp. 176-
In the
in inter
P. Jacqu

REVIEWS

Reports on Progress in Physics, Vol. XXIII.

Edited by A. C. Stickland. (The Physical Society, 1, Lowther Gardens, Prince Consort Road, London S.W. 7), 1960. Pp. 629. Price 63 sh.

The series of *Annual Reports on Progress in Physics* which the Physical Society, London, has been regularly publishing for the past nearly quarter of a century, have become an essential part of scientific literature in Physics. Nearly all colleges and centres of physical research possess these volumes, and each year they look forward to this annual addition to their libraries. A welcome departure was adopted in 1954 when the Society in addition to the publication of the bound volume also made available each article separately in paper covers at cheaper rates for wider use by individual workers interested in particular fields of study. Since the aims and objects of this publication as well as the general standard and quality of the articles in the annual issues are well known, we shall in this review of the latest issue Vol. XXIII, 1960, content ourselves with giving the titles of all the ten articles with their contributors and briefly indicate the importance of only some of them.

The first article is on "Optical properties of thin films" by O. S. Heavens (pp. 1-65). In this the author first discusses the optical behaviour of a surface layer (transition layer) of molecular dimensions as a problem in scattering, since the structure of such a film can be considered as aggregates of crystallites of variable size and orientation, and interspersed with different extents of voids. Multilayer films of several millimicrons thickness are treated from the electromagnetic theory, attributing to the layer a refractive index and (for absorbing material) an extinction coefficient. The article includes a discussion of interference filters, and some important applications of thin films in optics.

The second article is on "Group theory in solid state physics" by D. F. Johnston (pp. 66-153). The third is on "Photoelectric photometry" (pp. 154-75) by H. J. J. Braddick and the fourth article is on "Experimental analysis of the electronic structure of metals" (pp. 176-266) by A. B. Pippard.

In the fifth article on "New developments in interference spectroscopy" (pp. 267-312) P. Jacquinot gives a critical review of the

developments that have taken place in this field during the last ten to fifteen years. The article describes in particular the new types of spectrometers which select the different wavelengths by the amplitude modulation, and by the frequency modulation.

The sixth article is on "Planetary nebulae" by M. J. Seaton (pp. 313-54). Planetary nebulae are clouds of ionized gas surrounding certain hot stars. They have dimensions of the order of a million solar radii. Seen through a telescope they appear as pale-green discs resembling the planets Uranus and Neptune. The spectrum of a planetary nebula shows a large number of emission lines (both allowed and forbidden) against a background continuum. The pale-green appearance is due to the strong emission (forbidden) lines $\lambda 5007$ and $\lambda 4959$ of doubly ionized oxygen, originally ascribed to 'nebulium'. In this article the observed spectroscopic data are interpreted in terms of the processes taking place in a low density ionized gas exposed to dilute ultra-violet radiation from the central star.

The seventh article is on "Band structure calculations in solids" (pp. 355-94) by L. Pincherle.

The study of capture of an orbital electron by the nucleus is of fundamental theoretical importance as it points to the influence of the atomic electrons on nuclear properties. The most frequently observed capture is that of a K-electron, but capture of an L-electron can also occur. Orbital electron capture manifests itself only in the form of emission of X-rays and Auger-electrons. The article "Orbital electron capture by the nucleus" (pp. 395-452) by R. Bouchez and P. Depommier reviews the subject both from the theoretical and the experimental aspects.

In the article on "Precision measurement in γ -ray spectroscopy" (pp. 453-543) the authors G. A. Bartholomew, J. W. Knowles and G. E. Lee-Whiting discuss in detail high precision γ -ray spectrometers based on each of the four categories: (i) coherent scattering or crystal diffraction; (ii) photoelectric effect; (iii) Compton effect and (iv) pair production.

The last article is " ^3He induced reactions" by D. A. Bromley and E. Almqvist (pp. 544-629). On account of its high mass excess

(15-814 Mev.) and its high charge to mass ratio (2/3), the ^3He nuclide has come to be recognized as a very useful projectile in nuclear studies, and extensive publications have appeared in recent years dealing with ^3He induced reactions. The important results of investigations in this field have been summarized in this article. The report contains a comprehensive bibliography of papers covering about ten pages. The concluding part of the report containing suggested problems with low energy ^3He accelerators will be of particular value to research workers in the field.

A. S. G.

X-ray Spectrochemical Analysis. By L. S. Birks. (Interscience Publishers, Inc., New York), 1959. Pp. 137. Price \$ 5.75.

This is the eleventh volume in the series of monographs on Analytical Chemistry and its Applications. The X-ray spectrochemical analysis is not a new field of interest, but the development of new instrumental technique during the past decade necessitated the publication of a new book on the subject. This book under review is therefore a welcome addition to our shelves. As stated in the preface, the author has made an effort "to bring the subject up-to-date for the scientist who is interested in X-ray spectrochemistry as a research tool and also to present the material in a way that will be useful to those persons who are only interested in knowing enough about the methods to be able to use it judiciously for routine analysis". Some of the future potentialities have also been pointed out.

After a simplified but very short chapter (in 3 pages) on fundamentals, the author has arranged the subject-matter on excitation of the X-ray spectra, the dispersion and dispersion geometry of practical systems, and the detection and measurement of the spectra in the next three chapters. Following these are described techniques for quantitative analysis and their applications. In the seventh chapter the most important recent advance on X-ray spectroscopy, viz., the electron probe microanalyzer has been discussed in detail. This apparatus appears to have a great future because it can detect $\sim 10^{-13}$ to 10^{-14} gm. of an element or can perform chemical analysis on areas as small as 1 micron in diameter.

The general plan of the book was well conceived and has been written in a style to make the subject-matter readily comprehensible to the routine analysts and to newcomers. How-

ever, in a few places not very precise statements will be noticeable to a critical expert in the subject. Also, according to the reviewer, some discussion on X-ray absorption spectrography should have found some space in this book.

Though this book by Dr. Birks is not very comprehensive, it contains a considerable wealth of information on X-ray emission spectrography, provides useful suggestions to the worker and should prove to be a valuable addition to the literature of applied spectroscopy.

A. B. B.

Ultracentrifugation in Biochemistry. By Howard K. Schachman. (Academic Press, New York and London; India: Asia Publishing House, Bombay-1), 1959. Pp. 269. Price \$ 8.80.

No technique has done more than ultracentrifugation to the fundamental physical chemistry of macromolecules, and in recent years, it has become increasingly applicable to small molecules as well. As a tool for the preparation and separation of proteins, viruses, and many other substances it is of great and growing importance. Moreover, the processes occurring in sedimentation are complex and fascinating and the experimenter has therefore to understand them in order to interpret correctly the phenomena they observe.

In the present monograph are set out experimental and theoretical aspects of ultracentrifugation. After a brief introduction highlighting some of the principal developments, in Section II under the title 'General Considerations' the subject is dealt with in general terms. Section III deals with developments of the experimental techniques in the field, such as improvements in the instrument cells, rotors, measurement and control of temperature and the various optical systems. The following three sections deal with the fundamental principles of the major divisions, namely, sedimentation velocity, transient states and sedimentation equilibrium.

The presentation adopted in the monograph is likely to appeal more to the physical chemist than the biochemist, although the latter will get a grasp of the theoretical aspects of the subject by studying it. Details about experimental set-up are absent. A bibliography is appended at the end.

A. J.

Viscoe
Edito
New
Bom

The
civiliza
and car
plastics
resins
virtually
tries ha
modern

The
enginee
demand
proper
cations
strain
the o
approac
structur
not pos
perties,
cross-li
their m
factor
the bo
history

One
problem
menolog
sider th
which i
paramet
continua
mathemat
theories
the conc

These
in the ca
are still
the pub
giving i
theories
will be

The s
tents of
cussed a
was helo
ment Ce
Lancaste
and con
Analysis
(2) Th
Rubber-
pretation

Viscoelasticity—Phenomenological Aspects.

Edited by J. T. Bergen. (Academic Press, New York; India: Asia Publishing House, Bombay-1), 1960. Pp. x + 150. Price \$ 6.00.

The importance of polymers in life and modern civilization cannot be overemphasized. Proteins and carbohydrates are natural polymers. Rubber, plastics, synthetic elastomers and fibres, and resins are all polymers whose uses reach into virtually every facet of our lives. Their industries have assumed enormous proportions in the modern age.

The increasing use of polymer materials in engineering, design, equipment and construction demands a proper understanding of their elastic properties, at least in the range of their applications and production. The problem of stress-strain relations in polymers is different from the ordinary elastic problem. The basic approach is not a molecular approach but a structural one. In the case of polymers it is not possible to generalise their viscoelastic properties, since the presence of entanglements, cross-links, or crystallites profoundly affect their mechanical behaviour. Above all the time factor enters in the mathematical analysis and the boundary conditions are limited by the history of the process.

One method of theoretical approach to the problem which will be fruitful is the phenomenological approach, that is to say, to consider the viscoelastic material as a continuum which is characterized by specific, fundamental parameters, and to describe the response of this continuum to stress or strain by a system of mathematical statements applicable to the theories of continuum mechanics, and then test the conclusions from the results of observation.

These theories are less familiar and especially in the case of non-linear viscoelastic theories they are still in the formative stage. In this context the publication of the monograph under review giving in the compass of a handy volume latest theories and views on this important subject will be welcomed by workers in this field.

The seven chapters of this book are the contents of the seven papers presented and discussed at a symposium on viscoelasticity which was held in April 1958, at the Research Development Centre of the Armstrong Cork Company, Lancaster, Pennsylvania. The Chapter headings and contributors are as follows: (1) Stress Analysis for Viscoelastic Bodies by E. H. Lee; (2) The Linear Viscoelastic Behaviour of Rubber-like Polymers and its Molecular Interpretation by R. S. Marvin; (3) Comparisons of

Viscoelastic Behaviour in Seven Typical Polymer Systems by J. D. Ferry and K. Ninomiya; (4) The Behaviour of Certain Viscoelastic Materials in Laminar Shearing Motions by J. L. Ericksen; (5) Constitutive Equations for Classes of Deformations by R. S. Rivlin; (6) Stress Relaxation of Polymeric Material in Combined Torsion and Tension by J. T. Bergen; (7) The Normal Stress Effect in Polymer Solutions by H. Markovitz. A. S. G.

Lead Isotopes in Geology. By R. D. Russell and R. M. Farquhar. (Interscience Publishers, Inc., New York), 1960. viii + 243, Appendices I to XII. Price \$ 9.00.

The volume under review constitutes the first systematic attempt to bring together all the relevant data on the nature and geologic significance of isotopic abundances of common lead. The authors, Drs. Russell and Farquhar, have made original contributions to the subject and are hence eminently qualified to undertake such a compilation. Chapters I and II deal with the principles of lead isotopic method of dating and the techniques of determining lead isotopic abundances. Current ideas on the age of the earth, as deduced from lead isotopic data, are elucidated in Chapter III. The authors accept Patterson's value of $4,550 \pm 70$ m.y. for the age of the earth. The next Chapter relates to the principles of dating of galenas from their isotopic constitutions. Separate chapters have been devoted to anomalous lead, case histories, extension of Holmes-Houtermanns model (by Bate, Damon, Marshall and Russell) and lead-uranium-thorium methods of dating. Recent advances in the understanding of the significance of lead isotopic abundances have made it possible to decipher the geological history of a given region from a comprehensive study of lead isotopes. Such a study has been attempted for the regions of Sudbury and Thunder Bay, Canada and Broken Hill, Australia. The subject of rock lead, however, deserves a more extensive treatment than has been accorded to it in the present volume.

The twelve appendices, which appear at the end of the book, provide a complete list of lead isotopic analyses and thus constitute a valuable reference source.

The monograph is useful as a reference volume to researchers and post-graduate students interested in lead methods of dating.

U. A. N.

Proceedings of the Symposium on the Nature of Coal. (Central Fuel Research Institute, Jealgora, Bihar), 1959. Pp. 321.

The Central Fuel Research Institute, Jealgora, in collaboration with the Institute of Fuel (Indian Section) organised a symposium on the nature of coal and the volume under review consists of the texts of 40 papers presented at the symposium and the discussions thereupon. The symposium was organised in six Sessions, on the following topics: Origin and systematics of coal, Petrographic and X-ray studies on coal, Coal constitution—Physical methods, Coal constitution—Chemical methods and Physico-chemical properties of coal. It is interesting to note that commonwealth countries, U.S.S.R., Japan, etc., contributed about half the number of papers presented at the symposium.

The large volume of research that has been done on coal utilisation stands in marked contrast to the paucity of data on the nature of coal, its physical and chemical constitution and ultrafine structure. The present symposium highlights the recent advances made in India and other parts of the world on the nature of coal by the application of techniques like X-rays, electron microscopy, petrographic studies, etc., and suggests avenues of research which can be fruitfully explored.

The volume is a welcome addition to all research workers interested in fundamental research on coal.

C. MAHADEVAN.

The Chemistry and Biology of Sialic Acids and Related Substances. By A. Gottschalk. (Cambridge University Press, London N.W. 1), 1960. Pp. ix + 115. Price 22 sh. 6 d.

A great deal of interest has been evinced in recent years on the chemistry and biology of sialic acids since these compounds are known to provide the groupings which attract and bind influenza particles to the surface of host cells preliminary to the actual process of infection. The sialic acids are found widely distributed in animal tissues, glandular secretion and blood serum and form constituents of mucoproteins, mucolipids and lipoprotein-carbohydrate complexes.

After a historical introduction, Dr. Gottschalk has given an account of the physical and chemical properties of sialic acids as also their preparation from various natural materials, their quantitative assay and identification by specific colour reactions. He has then dealt with their distribution in tissues and body fluids under

physiological as well as pathological conditions. Further, there is a good account of Muramic acid which is the counterpart of sialic acid in bacteria and which appears to be an integral component of their basal cell-wall structure.

The author gives further in this monograph a well-balanced description of the molecular structures of sialic acids and related compounds as also an account of their known biological functions. As stated by him "it is the biological flavour which makes the cold beauty of the chemical structure displayed in this book so attractive" and it is to be hoped that a study of its contents will stimulate further research work in this new field of biochemistry.

P. S. SARMA.

Sector-Focused Cyclotrons—Proceedings of an Informal Conference held in Sea Island, Georgia, February 2-4, 1959. Edited by F. T. Howard. (Publication 656, National Academy of Sciences—National Research Council, Washington D.C.), 1959. Pp. 291. Price \$2.50.

The first cyclotron was built by E. O. Lawrence at Berkeley, and as a particle accelerator this device held supremacy for many years. Several artifices were introduced for obtaining greater and greater energy ranges. Relativistic range of energies were made accessible through frequency modulation. Another recent development is the use of sector focusing, to retain isochronism in the face of relativistic increase in particle mass. The basic difficulty with a cyclotron is, that the magnetic field should decrease with increasing radius to provide focusing, but should increase with radius to compensate for the relativistic mass increase of the particles during acceleration. L. H. Thomas first showed that this difficulty can be resolved by making the field to have azimuthal variations, which provide additional focusing to compensate for the defocusing. These azimuthal variations may be sinusoidal or square wave or radial or spiral. In the design, several factors enter such as choice of field shape representation, orbit calculations, adjustment of the field for the resonance condition to prevent phase slip, beam quality and design of radio-frequency system.

The forty odd papers presented during the Proceedings have fallen under six Sessions: Session I: Orbit Calculations and Magnetic Field Design; Session II: Realisation of Required Field Configurations and Model Magnet Work; Session III: Radio-frequency Systems; Session IV: Beam Quality; Session V: Beam Extraction;

Session VI: External Focusing Systems, Instrumentation, Operational Experience, and Summary of Outstanding Problems.

The Proceedings represent a considerable body of cyclotron operation experience, theory, design and testing work, informally presented, and is of much interest to experimental nuclear physicists.

A. J.

Chemical Analysis of Resin-Based Coating Materials. Edited by C. P. A. Kappelmeier. (Interscience Publishers, Inc., New York), 1959. Pp. 630. Price \$ 19.50.

Coating materials like varnishes and paints, compounded as they are with a base, a vehicle, a pigment and a thinner, are complex mixtures of inorganic and organic substances. With the increasing application of coating materials for industrial, protective and decorative purposes, and with the availability of many varieties of raw materials like oils, resins—natural and synthetic—and thinners, standardization and quality control of the raw materials and finished products demand a wide knowledge of analytical chemistry. The book under review supplies this demand. It is a valuable publication of great practical utility as it contains details of all the important methods used so far for the assay and analysis of coating materials. The editor, C. P. A. Kappelmeier, who had devoted a whole lifetime for the development of the paint industry, and twenty-six other specialists, have contributed the twenty-one chapters of the book. The subject-matter has been classified under three parts entitled: (I) Analysis of oil-based coating materials, (II) Analysis of lacquers and (III) Selected chapters on the analysis of coating materials. Part I contains methods for the separation of pigments and vehicles, as well as methods for the analysis of vehicles, oil-varnishes, resin-based coating materials, styrene-modified oils and alkyd resins, solvents and thinners, and latex paints. Part II deals with the analytical procedures for lacquers and resins. Application of infra-red spectrophotometry, steam distillation and complexometric titration and methods for the analysis of fatty acids, halogen compounds, silicone resins, polyester resins and isocyanate coating materials are included in Part III. The classical gravimetric and volumetric methods as well as the modern physical methods like spectrophotometry, polarography and chromatography and application in this field. The material presented is fully documented and indexed,

author- and subject-wise. A comprehensive list of items required to equip a laboratory for the analysis of coating materials is given as an appendix. It was unfortunate that the editor C. P. A. Kappelmeier did not live to witness the warm reception the book would receive at the hands of all those interested in the science and technology of coating materials.

B. H. IYER.

Principles of Dairy Science. By Ernest Vanstone and Bristow M. Dougall. (Cleaver-Hume Press Ltd., London W. 8), 1960. Pp. 238. Price 25 sh.

The scientific principles of dairying in relation to the nutrition of the animals, chemistry and bacteriology of milk and their application to processing for the fluid market and manufacture of milk products, legal aspects of processing and quality control, methods of quality control, water-supplies and sanitizers for the dairies, are covered in broad outline in the 21 chapters of the book. The text is presented in a lucid and logical sequence. Though there are a number of books on the subject, this new approach adopted by the authors in covering the field from a wider angle will appeal both to the students and busy executives as a source book for easy reference. The book has been clearly printed and well illustrated.

N. N. DASTUR.

Advances in Organic Chemistry—Methods and Results, Vol. I. Edited by R. A. Raphael, E. C. Taylor and H. Wynberg. (Interscience Publishers, New York), 1960. Pp. ix + 387. Price \$ 12.00.

In *Advances in Organic Chemistry: Methods and Results*, Volume I, has been reviewed (1) The Kolbe Electrolytic Synthesis; (2) Polyphosphoric Acid as a Reagent in Organic Chemistry; (3) Wittig Reaction; (4) Hydroxylation Methods; (5) The Selective Degradation of Proteins; (6) Optical Rotatory Dispersion and the Study of Organic Structures. As is evident the subjects comprise of methods used in synthetic and degradative studies, reagents with versatile application, and application of a very recent technique leading to precise and valuable information about conformation and configuration of organic molecules. References to latest developments and also to important unpublished results achieved in the field are of great advantage to the research workers. The introduction of "Experimental" section in different chapters is also very useful. The Editorial

Board should be congratulated on their success in getting the experts to write the different chapters.

D. K. BANERJEE.

Ticks, a Monograph of the Ixodoidea, Part V. By Don R. Arthur. (Cambridge University Press), 1960. Pp. xviii + 250. Price 60 sh. net.

The present volume forms the fifth part of a comprehensive work on the morphology, taxonomy, distribution, ecology and biology of ticks in relation to disease in man and other animals. The work itself was planned by Prof. G. H. F. Nuttall in the early years of this century when ticks were recognized as important carriers of *Piroplasma* and other parasites and the first part was published over 50 years ago. Dr. Arthur's book deals with five genera of which *Dermacentor* is the most important. Planned and executed with perfect care, the present part forms a fitting addition to the series on Ixodoidea and an invaluable reference work to all students of the group and also to parasitologists.

B. R. S.

***Aedes aegypti*.** By Sir R. Christophers. (Cambridge University Press), 1960. Pp. xii + 730. Price 75 sh. net.

Aedes aegypti is not only the celebrated yellow fever mosquito but also the insect most extensively used all over the world for research studies bearing on many general problems of the physiology and genetics of insects. Complete information on all aspects of the species is therefore not only desirable but also vital, wherever tropical diseases are being studied. Sir R. Christophers' work fills the need admirably. In virtue of his association with research on the mosquito for over 60 years, Sir Rickard has a comprehensive knowledge of the organism equalled by none. There is probably no work extant on the animal which Sir Rickard is not acquainted with and none of importance which he has not referred to in the preparation of this work, which will remain a classic for a long time. Every aspect of the mosquito is dealt with—systematics, structure, life-history, behaviour and physiology. Additional information on other mosquitoes is also provided and the work is an encyclopædia of information on the mosquito as a whole and *Aedes* in particular.

B. R. S.

Jute in India. By B. C. Kundu, K. C. Basu and P. B. Sarkar. (Published by the Indian Central Jute Committee, Calcutta. A Monograph), 1959. Pp. 395. 74 illustrations and charts and maps. Price Rs. 30.

This monograph is presented to the reader in three sections, with useful bibliography. Section I deals with the botanical, breeding and other aspects of jute crop. Jute is second to cotton amongst world's textiles. Of the 40 species known only *Corchorus capsularis* and *C. olitorius* are fibre producing. Excepting a few African, all Indian species are branching. The commercial jute being a bast fibre, its anatomy, physiology and the conditions determining the quality of the commercial fibre have been studied in detail.

Section II deals with the economics and analyses factors influencing the jute market. Nearly 80% of world's jute is produced in India and Pakistan and most world demand is met mainly by India. Brazil and Formosa also produce some jute. The most important export, apart from raw jute, is in burlap mainly to U.S.A. Serious growing competition is from European market exporting more highly finished burlap suited to the needs of U.S.A. To offset possible losses of markets exploration of new markets, developing internal ones, research to produce better finished products and to replace such imported articles are suggested.

Section III deals with the several technological aspects of jute fibre, interesting X-ray and chemical studies, processing, spinning, weaving and finishing. Species whose fibres are used as jute substitutes, those used in blending with jute, numerous possible uses of jute and other aspects are also dealt with. Though in many respects the technology is similar to that of cotton, in others it differs and creates new problems.

N. KRISHNASWAMY.

Cryogenics—An International Journal of Low Temperature Engineering and Research. Vol. I, No. 1, September 1960. Edited by K. Mendelssohn, R. B. Scott and L. Weil.

There is a need for a separate cryogenics Journal, especially on the engineering side, as the subject of cryogenic technology has in recent years advanced beyond the narrow confines of its applications to large-scale liquefaction and separation of gases only. With the increasing use of low temperature techniques in such widely different fields as liquid propellants, bubble chambers, physics of the solid state

chemist and engineering studies, imports, Journal, worker.

This wood consists of R. B. The large Abstracts the four and Russia.

The is of 5 "Cooling Nuclear distribution aspect research additional compilation a hundred and engineering during the

The a or \$15.0 by Heynations De Street, I

Books R Proceeding Congre by R. Press,

Introduc Advan Union, Pp. ix

Fourier (Stude bridge Pp. vii

The Major II. By versity xvi + 3 17 sh.

chemistry of free radicals, etc., the theoretical and engineering aspects of low temperature studies have come to occupy a position of importance in modern research. Hence this *Journal Cryogenics* will be widely welcomed by workers in this field.

This is a Quarterly Journal published by Heywood and Company Ltd., and edited by a Board consisting of K. Mendelssohn (Great Britain), R. B. Scott (U.S.A.) and L. Weil (France). The language of the Journal is English but the Abstracts of the original papers are printed in the four languages, English, French, German and Russian.

The first issue, Vol. I, No. 1, September 1960, is of 58 pages and contains a review article "Cooling by Adiabatic Demagnetization of Nuclear Spins" by N. Kurti, nine original contributions of about 4 pages each, covering different aspects of low temperature engineering and research, two letters, and two book reviews. In addition there is a bibliography which is a compilation giving the titles and authors of about a hundred papers on low temperature physics and engineering published in various journals during the period January-May 1960.

The annual subscription (4 issues) is £ 5, or \$ 15.00 or N.fr. 90. The Journal is published by Heywood and Co., Ltd., Scientific Publications Department, Carlton House, Great Queen Street, London W.C. 2, England.

Books Received

Proceedings of the Centenary and Bicentenary Congress of Biology, Singapore, 1958. Edited by R. D. Purchon. (University of Malaya Press, Singapore), 1960. Pp. 333. Price \$ 13.50.

Introduction to Physical Chemistry (Vol. III, Advanced). By S. N. Mukherjee. (Art Union, 80/15, Grey Street, Calcutta-6), 1960. Pp. ix + 832. Price Rs. 25-00.

Fourier Analysis and Generalised Functions (Students Edition). By Light Hill. (Cambridge University Press, London N.W. 1), 1960. Pp. viii + 79. Price 10 sh. 6 d.

The Major Achievements of Science, Vols. I and II. By A. E. E. Mackenzie. (Cambridge University Press, London N.W. 1), 1960. Pp. xvi + 368. Price 30 sh.; Pp. xi + 195. Price 17 sh. 6 d.

Cotton in India—A Monograph, Vol. II. By R. H. Dastur, R. D. Asana et al. (The Indian Central Cotton Committee, P.B. No. 1002, Bombay-1), 1960. Pp. viii + 339. Price Rs. 30-00.

Introduction to Submolecular Biology. By Albert Szent-Gyorgyi. (Academic Press, New York; India: Asia Publishing House, Bombay-1), 1960. Pp. 135. Price \$ 5.00.

Fluid Mechanics Through Worked Examples. By D. R. L. Smith. J. Houghton. (Cleaver Hume Press Ltd., 31, Wrights Lane, Kensington, London, W. 8), 1960. Pp. 344. Price 28 sh.

Harker's Petrology for Students. (Eighth Edition Revised.) By C. E. Tilley, S. R. Nockolds and M. Black. (Cambridge University Press, London N.W. 1), 1960. Pp. 283. Price 15 sh.

Physical Chemistry of Surfaces. By Arthur W. Adamson. (Interscience Publishers, N.Y.), 1960. Pp. xiv + 629. Price \$ 12.75.

Lectures in Theoretical Physics, Vol. II. Edited by Wesley E. Brittin and B. W. Downs. (Interscience Publishers, N.Y.), 1960. Pp. vii + 463. Price \$ 9.00.

Illustrated Genera of Wood Decay Fungi. By Charles L. Fergus. (Burgess Pub., Co., 426, South Sixth Street, Minneapolis-15, Minnesota), 1960. Pp. vi + 132. Price \$ 4.00.

Illustrated Genera of Imperfect Fungi. By H. L. Barnett. (Burgess Pub., Co., 426, South Sixth Street, Minneapolis-15, Minnesota), 1960. Pp. iii + 225. Price \$ 4.50.

Encyclopedia of Chemical Technology. Edited by Raymond E. Kirk and Donald F. Othmer. Second Supplement Volume, edited by Anthony Stonden. (Interscience Encyclopedia, Inc., New York-1, N.Y.), 1960. Pp. xv + 970. Price \$ 25.00.

Symposium on Monsoons of the World—New Delhi, 19-21, February 1958. Edited by S. Basu, K. R. Ramanathan, P. R. Pisharoty and U. K. Bose. (The Manager of Publications, Civil Lines, Delhi), 1960. Pp. x + 270. Price Rs. 12-00.

Laboratory Handbook of Toxic Agents. Edited by C. H. Gray. (The Royal Institute of Chemistry, 30, Russell Square, London W.C. 1), 1960. Pp. viii + 170. Price 20 sh.

SCIENCE NOTES AND NEWS

Incidence of Parasitism and Mortality in the Pupae of *Coccinella septempunctata* (Linn.)

Joginder Lall Nayar, School of Entomology, St. John's College, Agra, writes: More than a thousand pupae of *Coccinella septempunctata* (Linn.) were collected from the 'Brinjal' fields at Agra for the study of *Corpus luteum* formation in the said beetle. While rearing these at 84° F. and 40-45% R.H., it was observed that the total death of pupae amounted to 62.4%, out of which 50.7% were parasitised by the *Tetrastichus* sp. and 11.7% died due to other unknown reasons. This parasitism of the *Tetrastichus* sp. causing the destruction of the pupae of the predacious beetle is of great economic importance.

Award of Research Degrees

Madras University has awarded the Ph.D. Degree in Biochemistry to Shri S. Venkat Rao for his thesis entitled "Effect of Insect Infestation on the Chemical Composition and Nutritive Value of Foodgrains".

Andhra University has awarded the D.Sc. Degree in Physics to Shri M. S. V. Gopala Rao for his thesis entitled "Studies on the Spread of Irregularities in the Ionosphere" and the Ph.D. Degree in Physics to Shri S. Ramamurthy for his thesis entitled "The Application of the Theory of Molecular Orbitals to Certain Organic Molecules".

Utkal University has awarded the Ph.D. Degree in Mathematics to Shri Sibaprasad Misra for his thesis entitled "A Study of the Theory of Elementary Particles".

Symposium on 'Plant Embryology'

Under the auspices of the C.S.I.R. Biological Research Committee, a symposium on "Plant Embryology" was held at the University of Delhi from November 11-14, 1960. Professor P. Maheshwari, Professor of Botany at the University, was the convener of the symposium.

The symposium was attended by leading plant embryologists of the country and a number of young research workers. Thirty papers covering a wide ground were presented and discussed at the symposium. The following list gives the titles of some of the papers read: (1) Embryology of *Quinchamalium chilense* Lam., (2) Embryological studies in the Commelinaceae, (3) Embryological studies in the Loasaceae with special reference to endosperm haustoria, (4)

Embryological studies in relation to interspecific hybridization in jute, (5) *In vitro* studies on cotton ovules, (6) Intraovarian pollination in *Eschscholzia californica* Chem. and *Papaver rhoeas* L., (7) Embryology of *Paeonia* with a discussion on its systematic position, (8) The embryo of monocotyledons: a working hypothesis from a new approach, (9) The formation of male gametes in the pollen tubes of certain crop plants, (10) Megasporogenesis in duplication and deficiency heterozygotes of *Oenothera lamarckiana*.

There was also a discussion on "Teaching and research in plant embryology". Prof. P. Maheshwari suggested that a book on the "Comparative embryology of angiosperms" on the lines of Schnarf's "Vergleichende Embryologie der Angiospermen" be written as a joint undertaking by various authorities in India and abroad.—I. K. VASIL.

European Federation of Corrosion Annual Report for 1959

The Annual Report provides specialists with valuable information on the technical work on corrosion which has been carried out in Europe, with particulars of addresses and publications, research projects and papers read at meetings. The report can be ordered from the General Secretariat of the European Federation of Corrosion, Buro Frankfurt (Main), Postfach 7746. The price to members is DM 15 and to non-members DM 30.

Microbiology Congress to Meet at Montreal

The Eighth International Congress for Microbiology will be held at Montreal, Canada, from August 19 to 25, 1962, under the auspices of the Canadian Society of Microbiologists. There will be five Sections: Structure and Function; Agricultural Microbiology; Industrial Microbiology; Medical and Veterinary Microbiology; and Virology. Two or more symposia are being planned for each Section. All speakers at the symposia will be especially invited. There will be sessions for contributed papers in all Sections.

Enquiries should be made to Dr. N. E. Gibbons, Eighth International Congress for Microbiology, National Research Council, Ottawa-2, Canada.

United Nations Conference on New Sources of Energy

The United Nations Conference on New Sources of Energy, which will examine practical problems and experience in the utilization

of solar energy, from 21

The experts in energy participate in progress of view problems in these those are or facing

The Cations specific principle to or are applicat

to practice Requester Secretar Sources

Golden Instit

The I which o Year, h mark th have be of the I from 12 wide ra sciences ing, as t tions on applied nutrition engineer engaged has been weight current all reser

De novo

The b prophase attractive microscop brane or cence of reticulum the case units of

of solar energy, wind power and geothermal energy, will be held in Italy, probably in Rome, from 21 to 31 August, 1961.

The Conference aims at bringing together experts in these fields as well as those interested in energy development in general, to provide participants with up-to-date information on progress achieved and to facilitate an exchange of views and experience relating to practical problems, potentialities and limitations in utilizing these three sources of energy, especially in those areas lacking conventional energy sources or facing high energy costs.

The Conference will focus attention on applications rather than on the discussion of scientific principles and basic research, giving prominence to lines of action which have already led, or are about to lead, to commercial energy applications. Theoretical studies will be discussed only if they appear to be closely related to practical developments.

Requests for further information on the Conference should be addressed to: Executive Secretary, United Nations Conference on New Sources of Energy, United Nations, New York.

Golden Jubilee Research Volume of the Indian Institute of Science, Bangalore

The Indian Institute of Science, Bangalore, which observed 1959-60 as the Golden Jubilee Year, has brought out a Research Volume to mark the occasion. The papers in this volume have been contributed by the staff and students of the Institute. The 35 papers by 65 authors from 12 Departments of the Institute, cover a wide range of subjects in pure and applied sciences and engineering. These papers embodying, as they do, the results of recent investigations on diverse problems in chemistry, physics, applied mathematics, biochemistry, microbiology, nutrition, metallurgy, and different branches of engineering, will interest all research workers engaged in similar problems. The volume—which has been brought out in good print on heavy-weight paper, will be a useful addition to current scientific literature in the libraries of all research institutions.

De novo Origin of the Nuclear Membrane

The behaviour of the nuclear membrane in prophase and telophase is one of the most attractive features of mitosis for the electron microscopist and cytologist. The nuclear membrane or envelope is formed either by a coalescence of vesicles derived from the endoplasmic reticulum at the chromosome surface, or, as in the case of the onion cells, by slender lamellar units of the endoplasmic reticulum draping

themselves around the chromosomes at the poles in telophase. W. Bernhard has recently posed an intriguing question regarding the possibility of a *de novo* origin for the nuclear membrane.

In a problem like this, the selection of proper research material is perhaps more important than formulation of the original proposition. In such material, the number of cells undergoing mitosis should be unusually high, and the cells should not contain much endoplasmic reticulum and relatively few mitochondria. Electron microscope studies for an entirely different problem revealed that foetal liver (human and rat) is an excellent tissue for a number of cytological problems. At certain stages of development, the actively proliferating erythroblastic cells may outnumber the hepatic cells and even deform them by encroachment. In the youngest erythroblasts, fixed with buffered osmic acid, the sparse slender profiles of endoplasmic reticulum disappear gradually by fragmentation into small vesicles. Basophilic erythroblasts (normoblasts), which are characterized by an abundance of ribonucleoprotein particles and a small volume of endoplasmic reticulum or an involution of it, are mitotically quite active.

During prophase the nuclear membrane leaves the surface of the contracting chromatin, fragments and migrates towards the cell periphery. It is apparently metabolized after further fragmentation into vesicles. Although this lamellar structure may persist into the telophase stage, it is not related to the reconstitution of the new nuclear membrane.

Reconstitution of the nuclear membrane occurs first, multicentrically, around the periphery of the daughter cell chromosomal mass farthest away from the interzonal spindle fibres. Examples of aberrant chromosomal masses and differential mitosis further emphasize that remnants of lamellar endoplasmic reticulum are not required for the formation of the nuclear membrane in erythroblasts.

This evidence for the *de novo* origin of the nuclear membrane assigns to the chromosomal mass a much more important role in its formation than has been possible in previous electron microscope studies because basophilic erythroblasts are remarkably free from lamellar and vesicular endoplasmic reticulum. This interpretation is in keeping with light microscope studies showing that the nuclear membrane is first formed in telophase by the coalescence of membranes from the cytoplasmic surface of each of the chromosomal vesicles. Hence, the nuclear membrane stands in a direct genetic relation to the chromosomal parts, and in the case of basophilic erythroblasts, it would be

actively involved in the nucleo-cytoplasmic exchange of basophilic material—(*Nature*, 1960, 188, 239).

Electron Behaviour in Alloys

The problem of distribution of electrons in solid alloys consisting of one metal dissolved in another has attracted the attention of scientists in recent years. It has become increasingly clear that a proper understanding of the structure of alloys on an atomic scale is essential for the production of stronger materials on any but an empirical basis. The interactions between atoms in metallic solid solution can best be studied by the nuclear magnetic resonance method, where the resonance peak is profoundly modified by changes in the electron distribution round the solute atom. The application of this method has thrown new light on this problem. By observing the effect of alloying on the nuclear resonance of copper, T. J. Rowland of the Metals Research Laboratories of the Union Carbide Metals Company, has been able to determine how closely these extra electrons are grouped around the solute atoms. Observations show that the resonance amplitude of copper undergoes a sharp reduction upon alloying. Further, its dependence on the valency and size of the solute atom leads to the conclusion that the dominant source of the electric field gradients surrounding the solute atoms is the redistribution of the conduction electrons. Calculations based on new theoretical consideration have shown that the actual distribution of the electrons is as a diffuse cloud around the solute atoms. The charge density in the cloud decreases inversely as the cube of the distance from the solute atom rather than exponentially, and thus its effects are of longer range than had been supposed. It may be pointed out that earlier workers have often confused these electronic effects with "local strains" that also happen to decrease as the reciprocal cube of the distance from the solute atoms.

Visual Observations of Superconductivity Effects

An experimental technique developed in the G.E. Research Laboratory, Schenectady, makes

it possible to observe visually the transformations which take place in a superconducting material between the normal and the superconducting states. The method is based on two well-known facts connected with the phenomenon of superconductivity, namely, (1) a sufficiently strong magnetic field can change a material from the superconducting state to the normal state; and (2) so long as the material remains superconductive it is a perfect magnetic insulator.

In the experiment, the specimen to be observed is prepared in the form of a very thin flat disc. On top of this is placed a plate of a special cerium phosphate glass about $1/100$ " thick. When magnetized this plate has the property of rotating the plane of polarized light. A polarized beam of monochromatic light is made to fall on the glass, and the reflected light is viewed through a polarizing filter which can be rotated to any desired angle. The specimen material is cooled in liquid helium to a temperature of 1.5°K , when it becomes superconducting. But it can be changed back and forth between superconducting and normal states by varying a magnetic field. At certain magnetic field strengths the specimen is in the intermediate state with portions of it normal and portions superconducting.

Since the superconducting areas shield the portions of the glass plate directly above them from the magnetic field, these parts of the glass are not magnetized, whereas portions of glass directly above the normal areas, being not shielded, are magnetized. The magnetized areas of the glass rotate the plane of polarized light they reflect, while the shielded areas do not. By suitably adjusting the polarizing filter through which the reflected light is observed, the magnetized parts can be made to appear dark and the unmagnetized parts bright, or vice versa. Since the pattern of light and dark corresponds exactly to the pattern of superconducting and normal phases of the material, changes in the state of the sample can thus be visually observed or photographed—(*J. Frank Inst.*, 1960, 270, 250).

1007-60. Printed at The Bangalore Press, Bangalore City, by T. K. Balakrishnan, Superintendent, and Published by A. V. Telang, M.A., for the Current Science Association, Bangalore.

All material intended for publication and books for review should be addressed to the Editor, *Current Science*, Raman Research Institute, Bangalore-6.

Business correspondence, remittances, subscriptions, advertisements, exchange journals etc., should be addressed to the Manager, Current Science Association, Bangalore-6.

Subscription Rates: India: Rs. 12-00. Foreign: Rs. 16-00; £ 1-4-0; \$ 4.00.

FOR
**ACCURACY
AND
DEPENDABILITY**

in
CHEMICAL ANALYSIS & RESEARCH

use

laboratory reagents and guaranteed reagents

Manufactured by



SARABHAI MERCK PVT. LTD.

in conjunction with **E. MERCK A.G.** Darmstadt, Germany

Sarabhai Merck Pvt. Ltd., 13 Wittet Road, Bombay-1.

shilpi-s.m. 301

**SPECTROSCOPIC EQUIPMENT
AND
ACCESSORIES**

FOR
QUANTITATIVE ANALYSIS & RESEARCH WORK

Light sources like Arc Lamps with universal movements, precision slits, mountings for spectrographs, cameras, microphotometers, etc., etc.

Entirely Our Manufacture

For full particulars, please write to :

**THE GENERAL
ENGINEERING AND SCIENTIFIC CO.**

WALTAIR, VISAKHAPATNAM-3


(S. INDIA)

GRAMS : "ELECTRONIC"

Technical Adviser :

Dr. I. RAMAKRISHNA RAO

M.A., Ph.D. (Cal.), D.Sc. (Lond.)



INSIST ON "QUICO" PRODUCTS

"QUICO" STUDENTS MICROSCOPE

Stand of generous proportions inclinable by 90° with a fixed tube, without a condenser, with disc. diaphragm below the stage. Optical equipment: Nose-piece for 3 objectives; Originally imported German; 2 Achromatic objectives 10x & 42x; 2 Huygens eyepieces 10x and 15x; 4 different magnifications from 100 to 675x in a fine-wood cabinet with lock and key

Please Contact

Sole Selling Agents :

UNIQUE TRADING CORPORATION

221, Sheriff Devji Street


BOMBAY-3

PHONE : 30011

GRAMS : "UNILAB"



METRIC SYSTEM *in* CUSTOMS *and* CENTRAL EXCISE



The Metric System of Weights and Measures has come into force in Customs and Central Excise from October 1, 1960

Rates of duties are expressed in terms of metric units.

The rates have been converted to their nearest practicable equivalents in metric units.



METRIC SYSTEM

FOR SIMPLICITY & UNIFORMITY

Issued by Government of India

DA 60/39


The
ElectroniK
recorder
or indicator
is a
thousand
instruments
in one



An *ElectroniK* instrument adapts easily to your changing needs, never becomes obsolete. Its remarkable versatility is made possible by the many measuring circuits...many types of records or indications...many pen or print wheel speeds...and the wide variety of functions that can be incorporated in the instrument.

Use *ElectroniK* instruments to measure temperature, pressure, flow, pH, chemical concentration, voltage, speed — any variable that's translatable into a d-c signal.

Discover the advantages of *ElectroniK* versatility in specific applications...call in Blue Star to help you choose the right instrument for your application.



Honeywell

First in Control

Sold and serviced in India exclusively by

BLUE STAR

**BLUE STAR ENGINEERING
CO. (Bombay) Private LTD.**

LOTUS COURT

1 JAMSHEDJI TATA ROAD, BOMBAY 1

Also at CALCUTTA · DELHI · MADRAS

initiative from the shop floor


The manufacture of crowbars in Tata Steel's Agrico factory in Jamshedpur involved a number of time-consuming and uneconomic manual operations. Ratan Singh, a foreman, decided to do something to improve matters. Carefully studying the problem, he designed a die which cut down work, improved the product and substantially increased turnover at reduced cost. For his brilliant work, Ratan Singh received an award of Rs. 3,000.

In the ten years up to 1960, Tata Steel employees have put forward a total of 10,165 major and minor suggestions—an average of three a day. In the latest year, 445 suggestions were received, 54 of which have merited rewards.

Thus, initiative from the shop floor is helping to improve productivity at Jamshedpur, where industry is not merely a source of livelihood but a way of life.

JAMSHEDPUR

THE STEEL CITY



The Tata Iron and Steel Company Limited

TTN 4087

For
**ZOOLOGICAL SPECIMENS
 REQUIRED BY
 LABORATORIES
 AND
 INSTITUTIONS**



Consult:

Bombay Biological House

Dealers in Zoological Specimens

119, Hindu Colony, Dadar, Bombay 14

Established 1941

Phone : 61813

Gram : PHERETIMA

For
**EQUIPPING YOUR LABORATORIES
 IN**

**PHYSICS
 CHEMISTRY
 BIOLOGY
 GEOGRAPHY
 APPLIED MECHANICS,
 ETC.**

Contact

INDSALES Corporation

11, HORNIMAN CIRCLE
 BOMBAY-1

Phone : 253264

SP/GD/7.

BOROSIL LABORATORY GLASSWARE

such as

FLASKS, BEAKERS, CONDENSERS,
 MEASURING FLASKS, MEASURING
 CYLINDERS, PIPETTES & ANY
 SPECIAL APPARATUS MADE TO
 DESIGN

and


PENICILLIN VIALS, VACCINE BULBS—
 WHITE & AMBER



ALL OTHER APPARATUS & EQUIPMENT
 MANUFACTURED TO CLIENT'S DESIGN

**INDUSTRIAL & ENGINEERING
 APPARATUS CO. (PRIVATE) LTD.**
 CHOTANI ESTATES, PROCTOR ROAD
 GRANT ROAD, BOMBAY 7


TEMPO LABORATORY EQUIPMENT



(INCUBATOR Bacteriological)

TEMPO INDUSTRIAL CORPORATION (PRIVATE) LTD.

Sonary Road, Paranjpe 'B' Scheme, BOMBAY-57



pay for precision

BUY IRL PRODUCTS

**MICROSCOPES
and
PRECISION
OPTICAL
MEASURING
INSTRUMENTS**

India's Leading Manufacturers

Sole Selling Agents

MERCANTILE UNION PRIVATE LTD.

309, Bepin Behary Ganguly Street, Calcutta 12

Phone : 22-5778

**INSTRUMENT RESEARCH
LABORATORY LTD. CALCUTTA.**

NA IRL-3-42

**VERY RELIABLE INDIGENOUS SUBSTITUTES
OF GUARANTEED ANALYTICAL REAGENTS
MAY BE FOUND IN**

"BASYNTH"

**Brand
ANALYTICAL REAGENT**



Acid Hydrochloric
Acid Hydrochloric Fuming
Acid Sulphuric
Acid Nitric
Acid Nitric Fuming
Acid Acetic Glacial
Ammonium Hydroxide
Benzene
Toluene
Xylene
Petroleum Ether
Amyl Alcohol
Butyl Alcohol Etc., Etc.

Basic & Synthetic Chemicals (Private) Ltd.

P.O. Jadavpur University, Calcutta-32

CURR. SCI., DECEMBER 1960

MADE IN INDIA

**HIGH VACUUM
ROTARY PUMP**
SINGLE STAGE & TWO STAGE
with or without Air Ballast

All Indian materials and construction



**BASIC & SYNTHETIC CHEMICALS
(PRIVATE) LTD.**

P.O. Jadavpur University, CALCUTTA 32

Sigcol

LABORATORY
GLASS APPARATUS



Sole Selling Agents:

GHARPURE & CO.

P-36, ROYAL EXCHANGE PLACE EXTN.
CALCUTTA 1

Gram: "MEENAMO"

Phone: 22-2081



'STANDARD' RHEOSTATS

SINGLE, DOUBLE & FOUR-TUBE


COVER A WIDE RANGE
FOR USE IN

LABORATORIES & WORKSHOPS

Made by:

**THE STANDARD SCIENTIFIC
INSTRUMENTS CO.**

115, BRODIES ROAD, MADRAS 28



KEROY Short Beam Analytical Balance

No. K 1

Really Dependable Balance for Degree
Classes and Research Laboratories

Sensitiveness .. 1/10th mg.

Capacity .. 200 gm.

Catalogue on Request

Manufactured by:

Keroy (Private) Ltd.

BANARAS CANTT. :: CALCUTTA 10

"Calcutta Telephone No. is 24-3840"

Indian Journal of Dairy Science

Established in 1948 as a medium
for the advancement and dissemination
of knowledge in regard to dairy
science. Current Volume No. 13.

Editor: Dr. K. K. IYA

Subscription: Rs. 12
per volume of 4 parts, payable in advance

Single Copy: Rs. 3 each

*For advertisement rates and other
particulars, apply to*

**The Joint Secretaries
Indian Dairy Science Association
Hosur Road, Bangalore 1**

INJECTABLES — Safe and Dependable

TRADE



MARK

A wide range of parenteral preparations for meeting the growing requirements of the Medical Profession are now being processed in our Laboratories. They are made from Standard Chemicals employing double distilled and PYROGEN FREE water. Their containers (Ampoules) undergo rigid neutrality tests before they are selected for use. These Injectables are therefore guaranteed to be absolutely safe and dependable.

The following are but a few of our well-known Injectables :

- | | | |
|-----------------|---------|------------------------------|
| ● RETICULIN | | A Potent extract of Liver |
| ● HEXOPURIN | | An urinary Antiseptic |
| ● CALCITOL | | Injectable Calcium Gluconate |
| ● BEVITAMIN | | " Vitamin B ₁ |
| ● CEVITAMIN | | " Vitamin C |
| ● GLUCOSE SOLN. | | " Pure Dextrose |

The Mysore Industrial & Testing Laboratory Ltd.

Malleswaram P.O., Bangalore 3

Bengal Chemical and Pharmaceutical Works, Ltd.

The Largest Chemical Works in India

Manufacturers of Pharmaceutical Drugs, Indigenous Medicines, Perfumery, Toilet and Medicinal Soaps, Surgical Dressings, Sera and Vaccines, Disinfectants, Tar Products, Road Dressing Materials, etc.

Ether, Mineral Acids, Ammonia, Alum, Ferro-Alum, Aluminium Sulphate, Sulphate of Magnesium, Ferri Sulph., Caffeine and various other Pharmaceutical and Research Chemicals.

Surgical sterilizers, Distilled Water Stills, Operation Tables, Instrument Cabinets and other Hospital Accessories.

Chemical Balance, Scientific Apparatus for Laboratories and Schools and Colleges, Gas and Water Cocks for Laboratory use, Gas Plants, Laboratory Furniture and Fittings.

Fire Extinguishers, Printing Inks.

Office: 6, GANESH CHUNDER AVENUE, CALCUTTA-13

Factories: CALCUTTA - BOMBAY - KANPUR

*For the attention of Post-Graduate Teaching and Research
Laboratories*

**IF YOUR INTEREST
LIES IN**

**Botany }
Zoology }**

Chemistry

Medical Sciences

Physics

Geology

Research

YOU MUST HAVE

Bausch & Lomb 'STEREO-ZOOM'
Stereomicroscopes

Bausch & Lomb 'SPECTRONIC 505'
Automatic Recording Spectro-
photometer

Gallenkamp ADIABATIC Bomb
Calorimeter

Stanton UNIMATIC Single Pan
Balance

Bausch & Lomb 'SPECTRONIC 20'
Colorimeter—Spectrophotometer

Bausch & Lomb CERTIFIED-PRECI-
SION Diffraction Gratings (Replicas)

Ward's SYSTEMATIC Museum Collec-
tions

Gallenkamp HUMIDITY OVEN


Stanton Automatic THERMORECORD-
ING Balance

Please write to us for descriptive Catalogues as required

MARTIN & HARRIS (PRIVATE) LTD.

(SCIENTIFIC DIVISION)

SAVOY CHAMBERS, WALLACE STREET, BOMBAY 1



cenco

BETA RAY H/C METER

for

FASTER & ACCURATE MEASUREMENT

of

Percentage Hydrogen in Liquid Hydrocarbons

This instrument utilizes the absorption of beta rays as a means of measuring the percentage of hydrogen in the hydrogen-carbon ratio, in liquid hydrocarbons. It is a practical precision instrument designed for use in research as well as in product control and routine analysis. It is the fastest known means of obtaining important information, doing the entire job in a matter of minutes instead of the hours required by combustion methods.

The **Cenco Beta Ray H/C Meter** is accurate to within 0.02 weight per cent, hydrogen. Calibration is independent of pressure changes in ionization chambers since they share a common supply of gas. Temperature changes do not affect measurement since both specific gravity and beta absorption measurements are made automatically and at the same temperature. Adequate shielding makes the instrument safe from radiation hazards.

FULL DETAILS ARE SUPPLIED ON DEMAND

Sole Distributors

THE SCIENTIFIC INSTRUMENT COMPANY LTD.

ALLAHABAD

BOMBAY

CALCUTTA

MADRAS

NEW DELHI

Current Science, Vol. XXIX, 1960

Author Index

	PAGE		PAGE
A. B. B. (Rev.)	492	Bhargava, K. S.	443
Abdul Kareem, A.	413	Bhargava, P. D.	68
Abraham, A.	342	Bhaskara Rao, A.	16
Agarwal, J. P.	352	Bhat, J. V.	102, 181
Agarwala, S. C.	195	Bhatia, D. R.	24
Agnihotri, B. N.	285	—	116
Agnihotri, J. P.	352	Bhatia, I. P.	272
Agnihotrudu, V.	149	Bhatnagar, D. S.	193
Agarwal, H. S.	311	Bhatnagar, G. C.	483
Agrawal, N. S.	311	Bhatnagar, J. N.	391
—	311	Bhatnagar, M. P.	68, 187
Ajit Singh	134	Bhatnagar, S. P.	440
A. J. (Rev.) 71, 241, 287, 288, 327, 328, 407, 448, 449, 492, 494		Bhatt, Y. M.	439
Alice Abraham	100	Bhattacharjee, S. K.	48
Alikhan, M. A.	142	Bhattacharya, A. K.	128
Anantakrishnan, S. V.	51	Bhattacharya, H.	388
Anantha Samy, T. S.	133, 271	Bhattacharya, P. K.	147
Anikhindi, R. G.	90	Bhavnagary, H. M.	271
Annigeri, G. G.	319	Bhavsar, C. M.	177
Appala Raju, N.	52, 428	Bhawalkar, D. R.	175
Appa Rao, K.	23	Bheemasankara Rao, Ch.	136
A. R. V. (Rev.)	151	Bhowmick, K. K.	348
A. S. G. (Rev.) 111, 112, 245, 287, 367, 408, 409, 450, 491, 493		Bhujanga Rao, C. (Rev.)	370
A. S. (Rev.)	151	Bilgrami, K. S.	57, 227
Atal, C. K.	22, 56	Bilimoria, M. H.	181
Atheya, S. C.	353, 402, 403	Bir, S. S.	445
Aziz Khan, A.	135	Bora, K. C.	322
BABU RAO, P.	14, 467	Bora, P. C.	395
Badhe, N. N.	428	Bose, B. C.	391
Bal, D. V.	439, 489	Boszormenyi, Z.	340
Balakrishna, S.	476	Brewbaker, J. L.	298
Balakrishnan Nair, N.	61	B. R. S. (Rev.) 35, 114, 153, 155, 290, 496	
Balaram Sahoo	16	B. S. M. (Rev.)	336, 408
Ballal, D. K.	428	Butany, W. T.	141, 188
Banerjee, D. K. (Rev.)	495	CASSHYAP, S. M.	393
Banerjee, N. R.	223	C. C. P. (Rev.)	410
Banerji, S. N.	147	Chakravarty, H. L.	108
Basu, N. K.	136	Chandola, R. P.	187
Batra, H. N.	355	Chandy, K. C.	175
Beals, C. S.	205, 249	Chatterji, S.	108, 356, 479
		Chatterji, S. M.	219
		Chaturvedi, R. K.	128, 305

	PAGE		PAGE
Chaudhary, N. C.	193	Ganapathi, P. N.	273
Chavan, A. R.	32	Gandagule, B. N.	423
Chhibber, K. N.	320, 489	Gandhi, S. M.	42
Chopra, S. (Miss)	392	Ganguli, S.	44
Chopra, S. C.	108, 356	Ganguly, A. K.	30
Chopra, V. L.	362	Garga, R. P.	23
Choughuley, A. S. U.	93, 142	Gautam, O. P.	12
Choudhury, J. M.	64	G. B. (Rev.)	72, 23
Choudhury, K. D.	136	George, K. V.	11
Choudhury, S. K.	351	George, P. C.	31
Chughtai, I. D.	100	Ghose, K. B.	34
Cseh, Edith	340	Ghosh, R. B.	22
		Girdhari Lal	19, 47
DALBIR SINGH	489	G. N. R. (Rev.)	25
Das, N. B.	316	Godavari, C. P.	18
Das, S. M.	220	Gopala Krishna, G.	38
Das, Tam.	64	Gopalakrishnan, V.	24
Dastane, N. G.	134	Gopalan, C.	8
Dasannacharya, B. (Rev.)	111, 112	Gopal Rao, M. S. V.	34
Dasgupta, S. N.	226, 432	Govindu, H. C.	35
Dastur, N. N. (Rev.)	495	Guha Roy, A. R.	30
Dastur, R. H. (Rev.)	201	Guha, S. R. D.	8
Datar, M. G.	468	Gupta, B. D.	35
Datta, A. K.	181	Gupta, B. N.	43
Dave, J. S.	472	Gupta, P. D.	24
Davis, T. A.	273	Gupta, S. C.	146, 181
Deka, G. K.	292	Guram, M. S.	280
Deoras, P. J.	475		
Desai, M. C.	191	HALDANE, J. B. S.	375
Desai, M. V.	65	— (Rev.)	34
Desai, V. B.	476	Hanumantha Rao, M. S.	355
Deshmukh, G. S.	388	Hayakawa, M.	122
Deshmukh, M. J.	320	House, L. R.	295
Deshmukh, N. Y.	237	H. S. (Rev.)	114
Deshpande, B. B.	308		
Deshpande, D. D.	224	IDELMAN, S.	21
Deshpande, L. V.	385	Innes, M. J. S.	205, 240
Deshpande, P. K.	56	Irudayaswamy, A.	471
De Souza, E. J.	109	Iswaran, V.	63
Dharmatti, S. S.	165	Iyer, B. H. (Rev.)	495
Dhawan, N. L.	295	Iyer, V.	430
Dikshit, V. S.	484		
Divakaran, K.	487	JAGOTA, U. K.	437
Dominic, C. J.	274	Jain, B. D.	394
Dorai Raj, B. S.	277	Jain, J. S.	353, 402, 403
Doss, K. S. G.	472	Jain, R. C.	432
Dubey, A. S.	332	Jain, S. C.	345
Durve, V. S.	490	Jain, S. K.	361
Dutta, S. K.	388	Jaleel, S. A.	475
		Janardan Rao, Y.	17
ESHWAR, M. C.	388	Jande, S. S.	430
Ethirajan, A. S.	396, 400	Jauhari, O. S.	30, 69, 282, 430
		Jayaraj, R.	70
GADRE, G. T.	271	Jayaraj, S.	418
Ganapathi, K. (Rev.)	197, 201, 411	Jayaraman, N. (Rev.)	36
		Jha,	273
		Jogin	423
		John	42
		John	44
		Jone	30
		Jose	23
		Joshi	12
		Joshi	72, 23
		Joshi	11
		Joshi	31
		Juje	34
		J. V.	22
		KACH	375
		Kalia	18
		Kama	38
		Kane	24
		Kang	8
		Kano	34
		Kapo	35
		Kapp	30
		Kapu	8
		Karu	35
		Kar	43
		Kari	24
		Karta	146, 181
		Kaul	280
		Kaus	375
		Kaus	34
		Kelk	355
		Kemp	122
		Kish	295
		Kish	114
		Kohl	21
		Koth	205, 240
		Kova	471
		K. (.. ..	63
		Krist	495
		Krist	430
		Krist	437
		Krist	394
		K. R.	353, 402, 403
		K. S.	432
		Kum	345
		Kunc	361
		K. V.	475
		LAKS	17
		Laks	430
		Laks	30, 69, 282, 430
		Lal	70
		Lal	418
		Lall	36

	PAGE		PAGE
Jha, K. K.	231	Leela Devi, B.	99
Joginder Lal	441	Leelanandam, Ch.	225
Johnson, P. V.	476	Lewis, M. P.	52
John, V. T.	86	Lingaraj, D. S.	488
Joneja, M. G.	437	Loyal, D. S.	69
Joseph, J.	292	L. S. R. (Rev.)	36, 154
Joshi, A. B.	63, 285	Lukose, John	106
Joshi, B. S. (Rev.)	410	MAHADEVAN, C. (Rev.)	494
Joshi, J. M.	271	Mahadevan, V.	91
Joshi, R. D.	443	Majumdar, B. N.	434
Jujeja, S. L.	483	Malhotra, S. K.	282
J. V. B. (Rev.)	113, 244	Malik, C. P.	358
KACHRU, C. N.	55	Malik, Wahid, U.	135
Kaliaperumal, T. T.	485	Marimuthu, K. M.	482
Kamat, V. N.	133, 271	Mathai, A. O.	332
Kane, R. P.	425	Mathur, L. M.	116
Kanga, D. H.	430	Mathur, N. K.	92
Kanodia, K. C.	361	Mathur, R. S.	280, 353, 402, 403
Kapoor, J. N.	442	Mathur, S. C.	280, 353, 402
Kappanna, A. N.	196, 271	Mathur, S. P.	63
Kapur, O. P.	350	Meenakshi, Miss G.	438
Kapur, S. L. (Rev.)	328	Mehdiratta, R. C.	104
Kar, Amiya, B.	182	Mehta, C. M.	95, 431
Karibasappa, B. K.	326	Mehta, C. R.	95
Kartar Singh	24	Mehta, R. C.	95
Kaul, T. N.	31, 445	Merchant, J. R.	93, 142, 476
Kaushik, N. K.	316	Misra, G.	67, 105
Kaushik, R. L.	272	Misro, B.	480
Kelkar, V. B.	385	Mitra, S. N.	132
Kempanna, C.	138, 357	N. K. S. (Rev.)	154
Kishen Kumar	355	Mohanrao, G. J.	461
Kishen, K.	465	Mohapatra, K. C.	402
Kohli, V. P.	282	Mukerji, B. (Rev.)	152
Kothare, S. N.	109	Munjali, R. L.	442
Kovalsky, V. V.	171	Murty, G. S.	360
K. (Rev.)	328	Murty, Y. S.	158
Krishnamurti, B. G.	346	NAGARAJAN, G. R.	178
Krishnamurty, K. V.	23	Nagar, B. R.	270
Krishnan, M. S.	119, 161	Nag Raj, T. R.	192
— (Rev.)	196	Nair, K. N.	435
Krishna Rao, K. V.	386	Nair, K. S.	332
Krishnaswamy, N. (Rev.)	496	Nambiar, M. C.	234, 317
Kristensen, H. Rønde	124	Nanda, J. S.	281
K. R. (Rev.)	115, 200, 244, 369	Narain, A.	226
K. S. V. (Rev.)	150	Narasimhan, R. V.	76
Kumar, A.	195	Narayan, K. N. (Rev.)	290
Kundan Lal.	272	Narayana, L. L.	403
K. V. (Rev.)	448	Narayan, R.	472
LAKSHMANA RAO, M. V.	275	Narayan Rao, C. R. (Obituary)	173
Lakshmana Rao, S. K. (Rev.)	150	Naseeruddin Ahmad	185
Lakshminarayanaiah, N.	307	Natarajan, A. R.	471
Lal, B. M.	316	Natarajan, C. V.	461
Lall, B. S.	314	— (Rev.)	154, 197
		Natarajan, M. V.	393

		PAGE			PAGE
Nayar, B. K.	380	Purushottam, D.	308
Nayar, J. L.	372, 498			
Neelakantam, K.	52	QASIM, S. Z.	319
Nigam, R. K.	268			
Nirula, K. K.	489	RAI, J. N.	400
			Rajagopal, A. K.	13
COMMEN, C. N.	413	Ramachandran, K.	342
Oza, G. M.	25	Ramachandra Rao, B.	89, 304, 407	304
			Ramachandra Rao Pant, P.	304
PADHYE, M. R.	129, 344	Ramaiah, N. A.	306
Pancharatnam, S.	417, 457	Ramanathan, J. D.	131
Pande, B. P.	478	Raman, Sir, C. V.	1
Pande, P. G.	276, 302	Raman, V. S.	270
Pandey, N. D.	311	Rama Rao, H. S.	467
Pandya, H. G.	133, 271	Rama Rao, P.	140
Pandya, I. M.	268	Ramaswamy, M. K.	85, 335	335
Pandya, N. V.	14	Ramaswamy, M. N. (Rev.)	190
Panigrahi, B. M.	67	Ramdoss, S.	70
Panikkar, N. K. (Rev.)	88	Ram Gopal	483
Panikkar, N. K.	365	Ram Udhar	104
Panth, Bhola D. (Rev.)	10	Rana, N. G.	95
Pantulu, J. V.	28	Ranganathan, V.	330
Parameswar, N. S.	360	Ranga Setty, H. T.	246
Parthasarathy, S.	246	Rangaswami, G.	236, 318	318
Patel, A. R.	472	—	485
Patel, D. J.	247	— (Rev.)	330
Patel, G. H.	95	Rao, B. A. S.	473
Patel, I. D.	247	Rao, C. N. R.	52
Patel, N. K.	59	— (Rev.)	288
Pathak, C. H.	32	Rao, D. G.	350
Pathak, P. D.	14, 177	Rao, J. T.	321, 397
Patil, B. D.	63, 285	Rao, L. N.	190
Patil, B. V.	229	Rao, S. P.	92
Patnaik, D.	16	Rao, U. R.	425
Patnaik, S.	402	Rastogi, R. P.	268
Patra, B. D.	402	Rekhi, S. S.	237
Patwardhan, V. N. (Rev.)	199	Richharia, R. H.	480
Paul, R. C.	181	Rottenberg, J. A.	205, 240
P. B. (Rev.)	289	Roy, B. R.	132
Perur, N. G.	431	Roy, S. K.	189, 444
Perutz, M. F.	264			
Phatak, V. G.	25	SABNIS, S. D.	32
Pillai, S. C.	461	Sachdeva, S.	233
Pillai, S. K.	233	Sadasivan, T. S.	86
P. N. (Rev.)	33	— (Rev.)	112
Prabhakara Rao, A. V. S.	461	Sahu, D.	316
Prabhoo, N. R.	273	Saifi, A. Q.	391
Pradhan, V. G.	102	Saksena, K. N.	280
Prakash, U.	20	Salam, M. A.	148
Prakash Sarup	108, 356, 479	479	Salatore, S. A. (Rev.)	113, 155
Prasad, N.	66, 352	Sampath, S.	141
—	483	— (Rev.)	367, 288
Prasannan, K. G.	99	Sandhu, G. S.	222
Pritpal Singh	116	Sandhu, J. S.	222
Pruthi, J. S.	19, 473	Sandhu, S. S.	181, 222

	PAGE		PAGE
304	Sanghi, A. K.	187	Singh, R. N. (Rev.)
310	Sankaram, A.	97, 350	Singh, R. P.
	Santappa, M.	91	Singhal, S. P.
400	Santhanagopalan, T.	97	Sinna, A.
13	Saraswathy Royan	186	Sinha, P. C.
342	Sardar Singh	286	Sinna, U.
304, 467	Sarma, B. B. G.	145	Sirsi, M. (Rev.)
304	Sarma, P. S.	221	Sonalkar, V. G.
305	— (Rev.) 73, 74, 198, 243, 289, 369, 449, 494	494	Sondhi, M. M. (Rev.)
131	Sarwan Singh	268	Sood, S. M.
1	Sastry, C. A.	431	Sprague, E. W.
278	Sastry, G. R. K.	398	Sreekantath, G. M.
467	Sastry, M. P.	246	Sreeama Murty, K.
148	Satish Chandra	104	Srinivasachar, D.
85, 338	Satpanthi, P. S.	346, 429	Srinivasan, C. (Rev.)
190	Saxena, A. P.	90	Srinivasan, K.
70	Saxena, M. C.	128	Srinivasan, K. S.
483	Saxena, S. S.	281	Srinivasan, K. V.
104	Schwarting, A. E.	22	Srinivasan, M.
96	Seetharaman, R.	188	Srivastava, H. N.
393	Sehra, K. B.	183	Srivastava, O. K.
240	Sekariah, P. C.	276, 302	Subba Raman, P. R. (Rev.)
236, 318	—	230	Subbaram, M. R.
485	Sekhar, P. C.	230	Subba Rao, B. C.
330	Sen Gupta, P. N.	132	— (Rev.)
473	Sensarma, P.	108	Subba Rao, G. V.
52	Seshachar, B. R.	173	Subba Rao, K.
280	Seshadri, B.	21	Subba Rao, M. N. (Rev.)
359	Seshadri, T. N.	174, 433	Suolahmanyam, P. V. R.
21, 397	Seshadri, T. R.	53, 54, 131, 170, 178	Subrahmanyam, R. (Rev.)
190	Seshagiri Rao, D.	480	Subrahmanyam, V.
92	Seshaiya, R. V. (Rev.)	152, 412	Subramaniam, M. K.
425	Seshu, D. V.	326	— (Rev.)
268	Sethi, K. L.	360	Subramaniam, T. R.
237	Setty, T. H. V.	222	Subramanian, C. V. (Rev.)
480	Setu Rao, D.	51	Subramanyam, K. V.
5, 249	Shah, G. Z.	385	Subramoney, N.
132	Shah, H. M.	65	Suchir Kumar
39, 444	Shah, N. M.	131	Sulochana, C. B.
	Shama Kumari, K.	191	Sundara Rao, W. V. B.
32	Sharada, K.	130, 179	Susan Thomas
233	Sharma, C. P.	195, 437	Swaminathan, B.
96	Sharma, G. R.	219, 313	Swaminathan, M. S.
112	Sharma, H. N.	55	Synghal, K. N.
316	Sharma, P. D.	231	
391	Sharma, R. B.	60	TAIMNI, SHEELA
280	Shetty, B. V.	279	Tandon, K. K.
148	Shome, S. K.	432	Tandon, R. N.
3, 153	Shukla, R. R.	302	Tewari, A. P.
141	Siddiqi, R. H.	310	Tewari, J. P.
7, 286	Singh, B. D.	483	Thakar, G. P.
222	Singh, C. M. (Rev.)	370	Thakare, R. G.
222	Singh, K. P.	29	Thaker, B. M.
1, 222	Singh, L. M.	314	Thakur, M. K.
	Singh, R. D.	66, 484	Thakur, R. S.
			450, 451
			223
			390
			180, 225
			76
			231
		74, 154, 197, 291, 451	
			468
			363
			24
			295
			332
			14, 467
			364
			409
			321
			239
			76, 103
			350
			306
			470
			368
			86
			389
			368
			25
			89
			199
			461
			452
			307, 350
			186, 438, 482
			34
			76
			244
			55
			100
			229
			86
			63
			376
			246
			234, 298, 398
			139
			406
			62
			57, 227
			144
			400
			389
			322
			385
			138
			54

	PAGE		PAGE
Thangavelu, M.	99	Venkatakrishniah, N. S.	26
Thienemann, Prof. August (Obituary)	365	Venkataraman, T. V.	219, 313, 355, 441
Thombre, M. V.	191	Venkataraman, S. V.	324
Tiagi, B.	406	Venkatesan, R.	236
Tilak, S. T.	147	Venkatesan, V. K.	472
T. R. G. (Rev.)	329	Venkateswarlu, V.	126
Triloki Nath	270	Venkoba Rao, B. (Rev.)	451
Trivedi, J. M.	431	Venugopalan, P.	99
Trivedi, J. P.	131, 346, 429	Verma, S. C.	69, 280
		Verma, S. M.	345
U. A. N. (Rev.)	493	Vijayalakshmi, U.	103, 397, 400
Uma Agarwal	140, 284, 404	Vijayavargiya, R.	391
Umapathy, P.	428	Viladkar, B. G.	344
		Vishveshwara, S.	146
VAIDYA, P. C.	268	Vittal Rao, S.	454
Vaghani, Kum. D. D.	142	V. N. K. (Rev.)	73
Varadarajan, T. S.	129	V. (Rev.)	33, 71, 241, 287, 327, 366, 441
Varma, P. M.	359		
Vasantharajan, V. N.	236	WAHEED KHAN, A.	100
Vasil, I. K.	498	Wight, W.	395
Vasudeva, R. S.	359		
Vasudeva Menon, P. P.	413	YARAGUNTAIAH, R. C.	357
Vasudeva Murthy, A. R.	130, 179, 470	Yash Pal	41
Vedam, K.	79		

Subject Index

	PAGE		PAGE
<i>Acerentulus breviunguis</i> Conde from India	273	Amino-Acids in the Berries of <i>Withania somnifera</i> Dunal	22
Acoustic Thermometer	415	Amperometric Determination of Antimony and Thiocyanate	389
Adaptive Characters of Genetic Stalks of <i>Saccharum spontaneum</i>	321	Amphistome, Occurrence of Adult, in Sheep Lung	24
Additional Host, for <i>Korthalsella opuntia</i> (Thumb) Merr.	292	Analytical Applications of Tetrahydroxy Quinone	390
Additional Vectors of Tristeza Disease of Citrus	359	Anatase-Rutile Transformation, Impurity Effects on	52
Advanced Calculus (Rev.)	71	Anatomical Peculiarities of Two Clones of <i>Saccharum officinarum</i> L.	396
Advances in Enzymology (Rev.)	74	Animal Husbandry, Use of Tracer Atoms in	171
— in Inorganic Chemistry and Radio-Chemistry (Rev.)	151	Anomaly in Heat Capacity of Chromium	294
— in Organic Chemistry, Vol. I (Rev.)	495	Anthraxnose Disease of <i>Dioscorea alata</i> L.	66
— in Spectroscopy (Rev.)	328	Antibiotics Annual—1959-60 (Rev.)	411
<i>Aedes aegypti</i> (Rev.)	496	— in Medicine (Rev.)	451
Agricultural Fair, USA Exhibit, Co-60 Unit	298	— Therapy for Staphylococcal Diseases (Rev.)	197
— Institute, Allahabad, Golden Jubilee	332	Applications of Genetics to Cotton Improvement (Rev.)	201
— Society, Calcutta	202	<i>Araucaria bidwillii</i> Hook., Anatomy of Seedlings of	107
Aircraft Electrical Engineering (Rev.)	33		
Algebraic Theories (Rev.)	150		
Allyl Sulphide Content of Garlic	351		
Amino-Acid Constituents of the Carpet Beetle, <i>Anthrenus vorax</i>	479		

PAGE	PAGE	PAGE
Archegosaurus, a New Species of .. 144	Cell Cultures and Proteins .. 333	
Ashoka Guide (Rev.) .. 36	— Differentiation in the Regenerate of the Snail, <i>Helix aspersa</i> .. 478	
Asymmetrical Factorial Designs, On a Class of .. 465	— Organelles Stained with Acridine, Secondary Fluorescence of .. 186	
Atmospheric Radioactivity, Observations on .. 248	—, Organism and Milieu (Rev.) .. 244	
Auroral Green Line in Meteor Wakes .. 220	Centenary of Spectral Analysis .. 8	
Automatic Language Translator .. 414	Centromere in the Sex-Chromosome of the Males of <i>Hyalomma aegyptium</i> .. 437	
Award of Research Degrees: 37, 76, 116, 157, 246, 292, 332, 372, 413, 454, 498	<i>Cerataulus turgidus</i> Ehr. from Indian Waters .. 239	
Axenic Culture of Invertebrate Metazoa (Rev.) .. 35	<i>Cercospora</i> , New Species of, On <i>Leptadenia reticulata</i> .. 483	
Azo Dyes as Analytical Reagents for Aluminium and Beryllium .. 52	Cerous Nitrate and Potassium Salicylate, Complex Formation .. 128	
BACTERIAL Diseases of Sugarcane .. 318	Cervical Vertebrae of a Loris .. 355	
Balkan Lake Ohrid and its Living World (Rev.) .. 412	Cetyl Alcohol, X-Ray Diffraction of .. 175	
Barnacles from Bombay Shores, New Records .. 439	Chalkones, Reactions of .. 93	
Biochemical Society Symposium—Glutathione (Rev.) .. 330	Chemical Analysis: Analytical Chemistry of Titanium Metals and Compounds (Rev.) .. 242	
Biogeochemical Prospecting .. 456	— — of Air Pollutants (Rev.) .. 368	
Biogeography and Ecology in Australia (Rev.) .. 35	— — of Resin-Based Coating Materials (Rev.) .. 495	
Biology of the Amoeba (Rev.) .. 154	— — Surface-Active Agents (Rev.) .. 368	
— of the Pleuropneumonia-like Organisms (Rev.) .. 370	— Investigation of 'Oos' Natural Deposits Near Kapadvanj .. 131	
Bionomics of <i>Azygophleps scalaris</i> .. 311	— Pulp from Mesta Stick .. 348	
Biosynthesis and Secretion of Adrenocortical Steroid (Rev.) .. 369	Chemistry and Biology of Sialic Acids (Rev.) .. 494	
Black Bean Aphis, Response of, in Selecting Host-Plants .. 142	— of Heterocyclic Compounds s-Triazines and Derivatives (Rev.) .. 329	
Black-Tip Disease, Effect of, on the Catalase Activity of Mango Fruit .. 195	— of Natural Products, Vol. IV (Rev.) .. 410	
Blood Flow Rate, NMR Method of Study .. 77	Chlorine Filled Ozoniser as a Counter .. 90	
Blood Grouping and Human Trisomy .. 375	Chloropropamide and Diabetes Mellitus (Rev.) .. 74	
Blood-Groups (Rev.) .. 197	Chromatin Staining of <i>Coffea</i> Sp. with Basic Fuchsin .. 146	
Books Received: 36, 74, 115, 156, 201, 291, 331, 371, 412, 453, 497	Chromatogram, Method for Permanent Record of .. 21	
Building Research Workers' Conference .. 76	Chromatographic Behaviour of Cations .. 221	
CAESIUM Iodide, Thermal Expansion of .. 14	Chromosome Aberrations and Mutations in Nature .. 376	
<i>Casulia axillaris</i> Roxb., Morphology of .. 56	— Abnormalities in Man .. 374	
<i>Carissa Carandas</i> , Propagating by Air-layering with Growth Regulators .. 69	— Associations in <i>Gossypium barbadense</i> .. 398	
Carnegie Institution of Washington, Year-Book 1958 (Rev.) .. 407	— Complement and Meiosis in <i>Pteropodophus bimaculatus</i> .. 284	
<i>Carthamus tinctorius</i> , Colouring Matter of Flowers of .. 54	— Counts in Some Ferns .. 69	
Carvone Analysis by Ultraviolet Absorption in Indian Dill Oils .. 345	— Number and Sex Mechanism in Coleoptera .. 140	
Capsaicin Contents of Chilli Varieties .. 271	— — in <i>Leucas aspera</i> Spreng .. 231	
Cathode-Ray Tube and its Applications (Rev.) .. 328	— — of Some Himalayan Polytrichaceae .. 231	
	— —s in Some Species of <i>Vicia</i> .. 229	
	— —s in Two Compositae Weeds .. 360	
	— —s in Vitaceae .. 279	

	PAGE		PAGE
Chromosome Numbers of Dicotyledonous Plants	232	Ctenophores and Medusæ, Sudden Outburst of	382
— of <i>Berosus indicus</i>	404	Cultural Trials and Practices of Rice in India (Rev.)	206
— of <i>Oryza ridleyi</i> Hook.	326	<i>Curcuma longa</i> L., Flowering Behaviour and Anthesis of	402
<i>Cirrhina mrigala</i> , Absence of Pelvic Fins in	316	Current Virus Research (Rev.)	197
Citric Acid from Cotton	247	Cyanophyta (Rev.)	451
Clonal Propagation in Coconuts	273	<i>Cycas revoluta</i> , New Fungus on the Leaflets of	57
Clustering in Rice, A New Type of	188	Cytochemical Demonstration of Succinic Dehydrogenase	100
Cobalt-60 Irradiation Unit	298	Cytogenetic Investigations in Paniceæ	191
<i>Coccinella septempunctata</i> , Incidence of Parasitism and Mortality in the Pupæ of	498	Cytological Observations on <i>Asplenium</i> Linn.	445
Coconut Coir, High Stretch Paper from Coherence Properties of Electromagnetic Radiation—Part I	417	Cytology of Male Sterile <i>Crotalaria striata</i> D.C.	138
— — — — Part II	457	<i>Dacus zonatus</i> Saund, New Host Record of Darwin's Centenary Essays (Rev.)	372
Colchicine-Treated Phlox	281	Decay Scheme of Tl-210 (Rac")	455
<i>Colletotrichum</i> , a New Species from an Economic Host	147	Dechema Monographs (Rev.)	156
— <i>curvatum</i> on <i>Crotalaria juncea</i> , Occurrence of	357	Deep-Sea Fungi	333
<i>Colocasia</i> Embryos, Growing, in Culture	342	Dehydrogenase Activity in Plant Leaf Tissue, Use of 2, 3, 5-Triphenyl Tetrazolium Chloride	431
Colouring of Vanaspati with Curcumin from Turmeric	350	<i>Dendrophthæ falcata</i> (L.F.) Ettingsh, New Host of	158
Colour, Sensations of, and Visual Mechanism	1	Desert Locust, Occurrence of a Yellow Fledgling in	24
Complex of Bismuth Trichloride with Diethylamine	272	<i>Desmodium rotundifolium</i> DC., Bacterial Leafspot Disease of	65
— Formation between Cerous Nitrate and Potassium Salicylate	128	Detection of Copper by a New Spot Test	428
Condensation of Epichlorohydrin with Aromatic Amines	142	Diazinon, Colorimetric Method for Determination of	471
Conference on Building Materials	202	Diborane for Selective Reduction, Modified Method	389
— on Irreversibility and Statistical Mechanics	413	Digger Wasp Burrowing into Pith of Rose Plants	355
Confluent Hypergeometric Functions (Rev.)	287	Dipole Moment of 2-Nitro and 2-Bromo Para-Xylenes	51
Contemporary Physics (Rev.)	38	Discovery Reports (Rev.)	290
Continuous Spectrum of the Sun	218	Distribution of Pelagic Polychætes in the South Atlantic Ocean (Rev.)	452
Control of the Field Rat <i>Nesokia indica</i>	441	Dynamics of Particles and of Rigid, Elastic and Fluid Bodies (Rev.)	71
Copper Chlorophyll as Colouring Agent for Hydrogenated Vegetable Oil	349	EARTHQUAKE Shocks, Influence on Gravitometer Spring	476
— Murexide Complex, Spectrophotometric Investigation	128	Earthworms, Association between Two Adult	106
Coryneform Bacterium Producing γ -Carotene	102	Echo-I, the US Balloon Satellite	379
Cotton in India (Rev.)	451	Effect of Surgical Excisions of the Root Apex of <i>Sorghum vulgare</i>	233
— Seed Oil, Influence of <i>Aspergillus niger</i> and <i>Penicillium</i> on	100	Effects of Gamma-Rays on Growth of <i>Saccharum</i>	397
Coumaryl-Thioureas, Synthesis of	346		
Cruciferous and Non-Cruciferous Seeds, Microscopic Differentiation of	56		
Crustacean Wood-Borers from Andamans	275		
Cryogenics (Rev.)	496		
<i>Cryptostegia</i> , Studies in the Genus	32		

PAGE		PAGE		PAGE
392	Effect of Radiations on the Nutritive Quality of Bread Wheat	187	Field Resistance to Blight in <i>Colocasia antiquorum</i>	320
206	Elastic Behaviour of Matter under Very High Pressures	334	Flint \times Dent Hybrids in Maize	454
402	—, Dielectric and Piezo-Electric Constants of Quartz	333	Flora of Kutch, Additions to	361
197	Electrical Circuit Analysis (Rev.)	288	Flowers and Fruits on Root-Stock Stem of Mango Graft	285
451	Electrically Treated Cotton	117	Fluorescence Technique for Measuring Littoral Drift	373
57	Electrolytic Manufacture of Chemicals from Salt (Rev.)	196	Fluorescent Light Trap for Collection of Beetles	354
109	Electron Behaviour in Alloys	500	Fluoroxylenes, Near Ultra-Violet Absorption Spectra of	129
191	— Devices, Symposium on	170	Fluviological Deposits in the Outer Himalayas	104
445	— Emission from Silicon Carbide	38	Folic Acid, Purification of	388
138	— Paramagnetic Resonance—Two-Quantum Transitions	460	Formation of Complexes between Iodides	76
372	Electronic Nerve Cell	427	Fossil Meteorite Craters, Search for	205
384	— Pulse Technique and the Musical Scale	308	From Microphone to Ear (Rev.)	71
455	— Spectra of <i>o</i> -, <i>m</i> - and <i>p</i> -Chlorobenzaldehydes	344	Fuel Research in Australia	372
156	Electro-Osmosis in Charged Membranes	307	Fundamentals of Electronics (Rev.)	327
333	Elementary Treatise on Fourier's Series, Etc. (Rev.)	150	Fungal Pectolysis, Utilisation of	309
431	Elements of Radio Engineering (Rev.)	367	GEL-DIFFUSION Studies on <i>Toxoplasma gondii</i>	302
158	— of Wave Mechanics (Rev.)	33	General Cytochemical Methods (Rev.)	153
24	Embryology of <i>Eugenia malaccensis</i> Lam.	199	Genetic Basis of Selection (Rev.)	34
65	— of Staphyleaceae	403	Geochemistry of Rare and Dispersed Chemical Elements in Soils (Rev.)	409
428	Embryo-Sac Development in <i>Vernonia cinerascens</i>	406	Geology of the Country Around Polonnaruwa (Rev.)	196
471	Endotrophic Sporulation in <i>Bacillus cereus</i> , Inhibition of	430	Germfree Vertebrates: Present Status (Rev.)	35
389	Entomogenous Fungus (<i>Empusa</i> Sp.) Parasitic on Syrphid	219	Giant Mirror Telescope Designed	158
355	Entomological Conference	414	Gibberellic Acid, Effect on Germination of Pollen	332
51	Enzymes of Hepatics	104	— — — of, on Growth of Spinach	484
290	— of Two Common Dermatophytes	432	Gradientless Furnace	177
452	Enzymes (Rev.)	243	Gravimeter Spring, Influence of Earthquake Shocks on	476
71	Ergot in Bajra, Occurrence of	116	Gravitational Red Shift, Harwell Experiment	82
476	Estimation of Hydroxy Acids by Photochemical Oxidation with Ce (IV)	92	Gravitation and Electromagnetic Radiation	268
106	European Federation of Corrosion, Annual Report: 1959	498	Ground State of the C ₂ Molecule	9
379	<i>Euproctis howra</i> , Life-History of	230	Guide to Antibiotic Therapy (Rev.)	201
233	Excess Free Energies for Condensed Phases	268	Gyroscope Project, Very Low Temperature	373
397	Experiments and Exercises in Physical Sciences (Rev.)	196	HAEMATOXYLIN Squash Method for Root Tips of <i>Dolichos lablab</i>	482
	Eye Disease Causing Mortality of <i>Catla catla</i>	240	Harvey Lectures (1957-58) (Rev.)	154
	FADING of C.W. Signals as a Means of Spread-F Study	304	Heat and Thermodynamics (Rev.)	408
	Fallopian Tube, Organizations of Cells in	60	<i>Heliotropium indicum</i> , Control of, with Maleic Hydrazide	67
	Ferro- and Antiferroelectric Crystals	79	<i>Helminthostachys zeylanica</i> , Abnormalities in	190
	Ferromagnetics, Crystal-Oriented	40	Heterocyclic Chemistry (Rev.)	72
	Field Rat as Predator on Locust Hoppers	116		

	PAGE		PAGE
<i>Heterosporium</i> Blight of Onions, Occurrence of	31	Influence of Sex-Hormones on the Uptake of Zn ⁶⁵ by the Rat Liver	182
Heterotrichous Ciliate, <i>Metopus spiralis</i> , Occurrence of	435	Infra-Red and Raman Spectroscopy	263
High Altitude Fisheries with Special Reference to Kashmir	220	Inheritance of Leaf Chlorosis in Cross between <i>Triticum durum</i> and <i>T. dicoccum</i>	360
— Energy Accelerators, International Conference on	41	Inheritance of Leaf in Pigeon Pea	237
— Vacuum, High Temperature X-ray Camera	415	Insecticidal Trials against Termites Infesting Wheat Plants	108
History of Embryology (Rev.)	34	Insecticides for the Control of Sweet Potato Weevil	489
<i>Holcocera pulvereae</i> Meyr. as a Pest of Tamarind	413	Institute of Physics and the Physical Society	292
Holly Hock, Propagation of, by Air Layering	488	Institution of Chemists (India)	202
Hormones and Atherosclerosis (Rev.)	152	Interaction of Atoxyl with the Halogen Derivatives of Acetoacet Arylamides	431
Human Biochemical Genetics (Rev.)	289	— of Chromic Ions with Amino-Acids	135
— Gastro-Intestinal Tract (Rev.)	291	International Conference on High Energy Accelerators	41
— Integument, Normal and Abnormal (Rev.)	154	— on Nuclear Structure	335
— Mitotic Chromosomes, Nomenclature of	337	— Geodetic and Geophysical Union, XII General Assembly of	420
— Nutrition and Dietetics (Rev.)	199	— Oceanographic Expedition to the Indian Ocean	119
Hybrid Maize, Increased Food Production with	295	— Society of Tropical Ecology	76
Hydrated Compounds of Uranium from Uranium (IV) Oxycarbonate	15	Introduction to the Physics of Many-Body Systems (Rev.)	33
Hydrogenated Vegetable Oil, Copper Chlorophyll as Colouring Agent for	349	— to the Chemistry of Heterocyclic Compounds (Rev.)	448
Hypoglycemic Principles in the Roots of <i>Casearia esculenta</i> , Isolation of	138	Ionization Induced by Artificial Satellites	123
IAEA, Research Grants for Atom Scientists	157	Ion Thrust Device	40
Illustrated Genera of Rust Fungi (Rev.)	244	Irrigation Water, Economising by Biological Indicators	134
Incidence of Pink Bollworm on Cotton	440	Isolation of β - and γ -Sitosterols from <i>Clitoria marina</i> Linn.	225
Increased Food Production with Hybrid Maize	295	Isotopic Tracers (Rev.)	111
Indian Academy of Sciences	7	JEAN PICCARD, Award to	157
— Botanical Society	77	Joint British Committee for Vacuum Science	37
— Ephemeris for 1961 (Rev.)	408	Journal of the Marine Biological Association of India (Rev.)	114
— Institute of Science: Golden Jubilee Research Volume (Rev.)	499	— of Mathematical Analysis and Applications (Rev.)	37
Indian Science Congress	87	Jute in India (Rev.)	496
— Scientific and Technical Publications Exhibition (Rev.) 1960	287	KARRER Symposium on Vitamin A	157
— Society of Soil Science	76	LADY Tata Memorial Scholarships	37, 247
— Ocean, International Oceanographic Expedition	119	Lathyrism, Manganese and	86
Indole-Like Compounds in Parthenocarpic Guava	485	Lead Isotopes in Geology (Rev.)	493
Industrial Hygiene (Rev.)	10	Leaf Chlorosis, Inheritance of	360
Inflorescence of <i>Trichosanthes bracteata</i> Voigt.	108	Leucocyanidin-(Flav.-3-ene-3-ol)-Acetate, Preparation of	53
Influence of Cations on the Aggregating Effects of Synthetic Soil Conditioners	270		

- Leucocyanidin of the Groundnut, Constitution of .. 178
- Liquid Fluidised Bed Reactor .. 47
- Living Molecules .. 264
- Linear and Stereoregular Addition Polymers (Rev.) .. 328
- MACKERELS, Occurrence of .. 319
- Magnetic Annealing .. 294
- Sound Recording (Rev.) .. 71
- Magnetism and Molecular Structure .. 468
- Magnetohydrodynamic Generator .. 158
- Malachite Green as a Reversible Indicator in Acetyl Chloride .. 181
- Male-Sterile Off-Type Plants in *Sehima nervosum* .. 285
- Manganese and Lathyrism .. 86
- Mangiferin, Constitution of .. 131
- Masoniella indica* Sp. Nov. .. 148
- Mass Spectrometer, 100-Inch .. 117
- M-Chromosomes in a Pentatomid Bug .. 436
- Measurement of Power Spectra (Rev.) .. 366
- Mechanization of Thought Processes .. 70
- Medical Cyclotron for Studying Lung Function .. 121
- Meiosis in Pollen Sterile and Pollen Fertile Varieties of Coconut .. 234
- Melchior Treub (Rev.) .. 115
- Meliola albizziae* Hansford and Deighton from Assam .. 149
- Metallic Complexes of Embelin .. 136
- Metal Zone Plate to Focus Ultra-violet and Soft X-Rays .. 334
- Meteorite Craters, Search for Fossil .. 249
- Meteorological Rocket-Firing Network .. 382
- Meteor Trails as Reflectors of Radio Waves .. 160
- Methods of Biochemical Analysis (Rev.) .. 73
- Methylene-Bis-Derivatives, Formation of .. 95
- Mica in Cuddapah and Anantapur Districts .. 55
- Microbiology Congress .. 498
- Yesterday and Today (Rev.) .. 113
- Microcephalothrips brevipalpis*, Life-History of .. 437
- Microporous Plastic .. 455
- Microscope, a New 3-D .. 293
- Minor Constituents of Sea-Water .. 271
- Model Seismology .. 122
- Modern Co-ordination Chemistry (Rev.) .. 410
- Modern Fishing Gear of the World (Rev.) .. 83
- Molecular Complexes, in Gases, Detection by Ultrasonic Method .. 224
- Monazite in Granite-Gneisses .. 225
- Morazella caprae* Nov. Sp., Isolation of .. 276
- Morphology of the Endosperm in *Cesulia axillaris* Roxb. .. 56
- Mossbauer Effect .. 85
- Murexide as Indicator in Calcium Estimation .. 305
- Mylocerus viridanus*, a New Pest of Papaya .. 413
- NATIONAL Institute of Sciences of India .. 372, 455
- Physical Laboratory, Annual Report for 1959 (Rev.) .. 262
- Natural Hybrid between *Argemone mexicana* and *A. ochroleuca* .. 282
- Naturally Occurring Tetraploidy in *Nonnea pulla* .. 358
- Natural Purification of Flowing Sewage .. 461
- Rubber Research Conference .. 37
- Near Ultra-violet Absorption Spectra of Fluoroxylenes .. 129
- Necrosis in a Varietal Hybrid of Wheat .. 68
- Neutron Physics Symposia .. 415
- Radiation on Sex-Ratio in Watermelon, Influence of .. 322
- New Plant Fossils from Upper Gondwanas .. 145
- Nigrospora* Zimm., a New Species of .. 352
- Nitrification of Oiled and Deoiled Cakes .. 428
- Nitrogen Fixing Organisms in Coconut Retting Grounds .. 100
- Nobel Prize Awards for 1960 .. 422
- Nomenclature of Human Mitotic Chromosomes .. 337
- Nomenclature of Plants (Rev.) .. 290
- Nuclear Electronics (Rev.) .. 287
- Magnetic Resonance (Rev.) .. 407
- — —, Applications of .. 165
- Membrane, *De novo* Origin of .. 499
- Scattering of 765 Mev Nucleons, Total Cross-Sections for .. 385
- Science Series (Rev.) .. 407
- Structure, International Conference on .. 335
- Nucleoproteins (Rev.) .. 449
- OBITUARY: August Thienemann .. 365
- Obituary: Prof. C. R. Narayana Rao .. 173
- Occurrence of Mackerels Off Ratnagiri Coast .. 319
- Oceanographic Expedition to the Indian Ocean .. 119
- Oleaceae, Floral Structure of .. 59
- Olpidium uredinis* Parasitic within Urediospore of *Uromyces leptodermus* .. 229
- Onion Powder, Equilibrium Relative Humidity .. 19
- Smut in Mysore .. 324
- Thrips, On the Biology of .. 314

	PAGE		PAGE
<i>Ophioglossum</i> from the Plains of Eastern U.P.	444	Physical Methods of Investigating Textiles (Rev.)	242
Ophthalmology, Academy of, First Congress	157	— of Organic Chemistry (Rev.)	367
Optics and Spectroscopy (Rev.)	78	— Research Committee, Conference	83
Oocystis (<i>O. kumaonensis</i> Sp. Nov.), a New Species	29	Physiological Studies on the Cotton Crop (Rev.)	244
Organic Insecticides, Effect of, on Crop Yield—Part II	480	Physiologic Races of <i>Sphacelotheca sorghi</i>	226
Oriented Graphite	159	Pinguite, Remarks on	16
Organic Chemistry (Rev.)	72	Pioneer V, the U.S. Sun Satellite	116
— (Rev.)	330	Pistil-less and Functional Male-Sterile Plants	63
— Insecticides against Termite Damage in Cotton Crop	356	Plant Resources, UNESCO Symposium on	454
Origin of the Blue Continuum in the Hydrogen Flame	247	— Virus Diseases in Denmark	124
— of the Solar System	343	Plastic Capillaries for X-Ray Diffraction	333
Orthogonal Polynomials	13	Pollen Irradiation and Culture of Pollen Tubes of Coconut	317
<i>Oryza</i> Species, Response to Photoperiods of	141	Pollution of the River Kali by the Effluents of Industrial Wastes	310
Ovule Development in Carpelloid Stamens of a <i>Nicotiana</i> Hybrid	23	<i>Polydora</i> as an Agent in the Destruction of Submerged Timber	61
Oxalate Content of Leafy Vegetables	133	Potential Constants of Certain Planar XY ₂ Z Molecules	487
Oxidation-Reduction Potentials of Organic Systems (Rev.)	288	— of VOCl ₃	14
Palaeobotanical Society, Thirteenth Annual Meeting	413	Powder Chromatography for Carotene Estimation	434
Palm Fruits from the Deccan Inter-trappean Beds	20	Prawn Larvæ, New Technique for Preserving	238
Paper Chromatography of Antioxidants — Electrophoresis, Simple Apparatus for	97	Pre-meiotic Somatic-Reduction in Wheat Preservation of Fruits and Vegetables (Rev.)	362
— from Coconut Coir	93	Principles of Dairy Science (Rev.)	370
<i>Parasa lepida</i> Cram. on Arabica Coffee	246	— of Quantum Electrodynamics (Rev.)	495
Pattern of Food and Blood Cholesterol	183	Probability and Related Topics in Physical Sciences (Rev.)	241
Pectin Decomposition by Actinomycetes	181	Proceedings of Symposium on Nature of Coal (Rev.)	327
Pelvic Fins, Absence of, in <i>Cirrhitina mrigala</i>	316	— of the Fourth Congress on Theoretical and Applied Mechanics (Rev.)	494
<i>Pennisetum typhoides</i> , Accessory Chromosomes in	28	Processed Plant Protein Foodstuffs (Rev.)	150
Perfect Stage of Apple Scab Pathogen, Occurrence of	445	Propagation of Peach by Stem-Cuttings	151
— of <i>Phyllosticta cycadina</i>	227	Proportionality Errors, Demonstration of	282
Perfumes, Cosmetics and Soaps (Rev.)	199	Protactinium Stock	364
Petrology of the Clastic Dykes of East Bokaro Coalfield	393	Protein from Breathed-in Nitrogen	118
Pharmacological Actions of <i>Visna rosea</i> Linn.	391	Proteins, Symposium on	172
Phenols, Some Reactions of	476	Proton Microscope	247
Photoperiodism in Plants and Animals (Rev.)	369	Protopectinase Enzyme Secretion by <i>Penicillium expansum</i>	77
Photo-Piezoelectric Effect in Semiconductors	116	Pteridaceæ, Studies in	146
Photosensitized Polymerisation by Uranyl Ions	91	QUANTUM Chemistry (Rev.)	380
		Quartz Veins in the Yellandlapad Rocks	448
		Quinaldine, Use of, in Transporting Live Fish	17
			393

PAGE		PAGE		PAGE
	RADIATION Belt Around Jupiter ..	293	Scattering of High-Energy Electrons by	
	— —s, Study by Nuclear Emulsion ..	159	Carbon and Oxygen ..	385
242	Radio Galaxies, New Hypothesis on For-		Scientific Papers of Sir Geoffrey Ingram	
367	mation ..	38	Taylor (Rev.) ..	366
83	Radioisotopes Aid Indian Agricultural		Search for Fossil Meteorite Craters—I ..	205
	Research ..	46	— — — — II ..	249
244	Raman Spectra Investigations, Differential		Sea-Water, Minor Constituents of ..	271
226	Method for ..	84	Secretory Activity of the Funnel of Avian	
16	— Spectroscopy, Infra-Red and ..	263	Oviduct ..	274
116	Raptakos Medical Research Fellowships	37,	Sector-Focussed Cyclotrons (Rev.) ..	494
		475	Sectorial Chimera in Onion ..	483
	Rats, Studies on Bombay ..	475	Selaroides leptolepis, Food and Feeding	
63	Rauwolfia canescens Linn., Occurrence in		Habits of ..	62
454	Ganjam District ..	25	Semiconductors of Organic Polymers ..	39
124	Reaction between Piperidine and Carbon		— (Rev.) ..	448
333	Tetrachloride ..	179	Semimicro Experiments in General	
	— Products of Zirconium Tetrachloride		Chemistry (Rev.) ..	196
317	with Phenols ..	222	Seminar in Electrochemistry ..	372
	Recent Progress in the Endocrinology of		Sensations of Colour and the Nature of	
310	Reproduction (Rev.) ..	152	the Visual Mechanism ..	1
	Redox Processes, Symposium on ..	247	Separation of Uranium and Rare Earths	308
61	Red Shift, Gravitational, Harwell Experi-		Sewage Natural Purification of ..	461
	ment to Measure ..	82	Shell Disease in <i>Crassostrea gryphoides</i>	
487	Red-Spider Resistance in Tea ..	395	(Schlotheim) ..	489
14	Reflex Klystrons (Rev.) ..	241	Sitosterol from Seeds of <i>Clitoria ternatea</i>	180
	Reflector Telescope of the Tautenburg		Sky Component Equations for Overcast	
434	Observatory ..	424	Sky ..	174
	Regeneration of Erythrocytes, Effect of		Sky-Component Surfaces ..	433
238	Injected Methionine on ..	99	Smut on Onion and Garlic ..	26
362	Relationship between Milk Production and		Soil Fungi, New Records in ..	400
	Body Measurements in Murrah Buffaloes	193	— Humidity by Radioactive Methods ..	127
	Relaxation Times of Dipolar Liquid		— Salinity, UNESCO Training Course on	454
370	Mixtures at 3 cm. ..	306	Solar Radiation Pressure on Satellites	421
495	Reports on Progress in Physics, Vol. XXIII		— Spectra from Rocket Flights ..	117
241	(Rev.) ..	491	— X-Rays ..	159
	Reproduction in Domestic Animals (Rev.)	289	Solid State Physics, Symposium on ..	83
327	Resistance of Barley Varieties against		Solvent Effects on the Displacement of	
	Smut ..	280	Raman and Anti-Stokes Lines ..	303
494	— of Gram Varieties to <i>Fusarium</i> Wilt	403	— Extraction of Vegetable Oils (Rev.)	113
	Resorpiophenone Oxime as a reagent for		Source Book in Mathematics (Rev.) ..	448
150	Copper Determination ..	472	Spectral Analysis, Centenary of ..	8
151	<i>Rhinoceros unicornis</i> ..	77	Spectrophotometric Investigation on	
282	Rhizobia in Inoculated Seed, Survival of	350	Copper Murexide Complex ..	128
364	Root-Knot Nematodes on Sugarcane ..	236	Sputnik—III Disintegrates ..	203
118	Ruby Maser ..	334	Standard Frequency and Time Transmis-	
172	Russian Journal of Physical Chemistry		sion Centre in Delhi ..	158
247	(Rev.) ..	202	Sterility in Rice, Influence of Day Length	
77	— Space-Ship—II ..	383	on ..	105
	<i>Saccharum officinarum</i> , Pattern of Pubes-		Structure Reports (Rev.) ..	288
146	cent Culms in ..	400	Studies in Medical Mycology ..	432
380	— — Anatomical Peculiarities ..	396	— in Theoretical Physics (Rev.) ..	336
448	<i>Sarcopodium</i> , a New Species of, on Coffee	192	— on Bombay Rats ..	475
17	Satellite, Soviet ..	294	— on Structure and Development of	
	Scale Insect on Betel Vine ..	76	Vertebrates (Rev.) ..	154

	PAGE		PAGE
Sugarcane Mite <i>Oligonychus indicus</i> , control of	313	Tillering in <i>Sorghum</i> , Factors for ..	487
Sulfur in Proteins (Rev.)	198	Titrimetric Determination of Copper ..	223
Sulphur Dioxide and Piperidine Complexes	130	Tracer Atoms, Use of, in Animal Husbandry	171
Summer School in Botany	301	<i>Thorea ramosissima</i> Bory in India, Occurrence of	490
— of Theoretical Physics	293	Tracheal System in Agromyzid Larvæ ..	311
Sun as Relay for Radio Signals	373	Translocation of Hydrolysed Substances from Endosperm to Embryo, Effect of Light on	64
Survival of Rhizobia in Peat-Based Legume Inoculants	97	Treatise on Algebraic Plane Curves (Rev.) ..	150
Symposia of the Society for Experimental Biology (Rev.)	112	Triboluminescence in Mercury Bubbler ..	373
— on "Advancing Frontiers of Life Science"	332	Triploidy in Ramie	191
— on Light Metal Industry in India ..	372	Trisomy, Blood Grouping and Human ..	375
— on Minor Batch Constituents in Glass Melting	117	Tritium in Nuclear Fission	456
— on Plant Embryology	498	Tungus Meteorite, New Hypothesis on ..	414
— on "Proteins"	247	Tunnel Diode	39
— on "Redox Processes"	247	Turbidimetric Chloride Determination ..	130
Synthesis of (2-Alkyl-4:5-Dimethoxy Phenyl)- <i>n</i> -Propylamine	55	Two-Parameter Discrete Approximation to any Unimodal Continuous Distribution ..	246
— of Chlorosubstituted Chalkones	429	Two-Quantum Transitions in Electron Paramagnetic Resonance	460
Synthetic Diamonds	38	ULTRACENTRIFUGATION in Biochemistry (Rev.)	492
TANDEM Satellites in <i>Cicer arietinum</i> Linn.	438	Ultrasonic Absorption and Velocity in Water Containing Algae	333
<i>Tanymericus</i> Weevil in Pest Form	286	— in Some Pure Liquids	467
Tape-recorder with 40 Hours Continuous Reproduction	455	Ultrasonically Dispersed Sodium	50
Tapioca as a Source of Alcohol	346	Ultrasonic Flowmeters	203
Technetium	160	— Method of Detection of Molecular Complexes in Gases	224
Tectonic History of the Earth, Two Stages in the	203	— Studies in Glycerol-Alcohol Mixtures ..	89
— Pattern of India	161	Ultra-Violet Radiometry Standard	160
Television and the Electron Microscope ..	117	Underwater Sound by <i>Therapon jarbua</i> Production of	277
Termites, Symposium on	292, 454	United Nations Conference on New Sources of Energy	497
<i>Terminalia crenulata</i> Roth., 4-Winged Fruit of	25	University Physics (Rev.)	450
Test-Tube Protein	78	Unrecorded Diseases of <i>Sorghum</i> and Maize from India	442
Tetrahydroxy Quinone, Analytical Applications of	390	Uptake and Reduction of Iodate by Wheat-Roots	340
Text-Book of Physics (Rev.)	111	VANADYL Sulphate and Malonic Acid, Reaction between	147
Theoretical and Applied Mechanics Sixth Congress	413	Vanaspatis, Colouring with Curcumin from Turmeric	350
Theory of Numbers and Diophantine Analysis (Rev.)	150	— Industry (Rev.)	155
Thermal Expansion of Cæsium Iodide ..	14	—, Vitamin K as Colouring Agent for (see Also Hydrogenated Vegetable Oil) ..	221
Thermogravimetric Behaviour of Lanthanum, Thorium and Cerium Oxalates ..	470	Varennas Summer School on Weak Interactions	40
Thiamine and Biotin, Requirements of <i>Glomerella tucumanensis</i>	103	Varietal Reaction of Jowar to Grain Smut	463
Ticks, a Monograph of the Izodoidea (Part V) (Rev.)	496		

PAGE		PAGE		PAGE
487	Varietal Resistance of Wheat to Loose		Window in the Sky (Rev.)	403
223	Smut	353	Wind Tunnel for High Speed in	
	Vegetative Propagation in Rice Breeding	480	Amsterdam	118
171	— of <i>Hevea brasiliensis</i> by Cuttings ..	414	World Agricultural Fair, Co-60 Unit ..	298
	Vertebrate Fossil-Bone from Umia Beds ..	185		
496	— Story (Rev.)	155	X-RAY Diffraction of Cetyl Alcohol,	
311	"Vibrated" Dropping Mercury Electrode ..	472	Effect of Ageing and Thermal History	
	Virus Disease of Plants in Madhya		on	175
	Pradesh	324	— Method for Determining Grain-Size in	
64	— Growth and Variation (Rev.)	73	Rocks	386
150	Viscoelasticity (Rev.)	493	— Precession Photographs in three Dimen-	
373	Visual Observations of Superconductivity		sions, Simple Device for Viewing ..	456
191	Effects	500	— Spectrochemical Analysis (Rev.) ..	492
375	— Pigment of the Horseshoe Crab	248		
456	Vitamin B ₁₂ , Role of, in Nitrogen Fixation	63	YEAST, Heat-Resistant and Acid-Tolerant,	
414	— K as Colouring Agent for Vanaspati ..	221	from <i>Passiflora edulis</i>	473
39				
139	WATER Discharge of World Rivers	247	<i>Zanthoxylum rhetsa</i> , DC., Investigations	
	Water-melon Mosaic Virus	443	on	95
246	Watumull Memorial Awards	292	Zeeman Effect, a Simple Experimental	
	Wave Propagation, Symposium on	170	Arrangement to Study	332
460	Weak Interactions, Varenna Summer		Zenith Angle Response for Inclined	
	School	48	Meson Telescopes	425
	Wealth of India: Industrial Products—		Zinc Status of Indian Soils	316
492	Part IV (F-H) (Rev.)	245	<i>Ziziphus mauritiana</i> , Lam. (Ber), Vege-	
	—: Industrial Products—Part V (I-L)		tative Propagation of	30
333	(Rev.)	409	Zoogeography (Rev.)	114
467	—: Raw Materials—Vol. V. (H-K)		Zygnemaceæ (Rev.)	450
50	(Rev.)	112		
203				
224				
89				
160				
277				
497				
450				
442				
340				
147				
350				
155				
221				
40				
403				

